

A Framework for Assessing Heat Risk and Adaptation Capacity in Local Communities

Sr. No.	Author Name
1.	Ms. Drashti Patel
2.	Ms. Prerna Chavan
3.	Ms. Mansi Koli
4.	Ms. Kalyani Wable

Guide: - Mrs. Anjali Dandekar anjali.dandekar@ruparel.edu

Assistant Professor

Modern Education Society's "D. G. Ruparel College of Arts, Science and Commerce" Matunga West

Abstract: Heat waves and rising average temperatures are among the most direct and immediate impacts of climate change. This paper examines how heat stress affects communities, presents a simple method to assess local risk, and evaluates low-cost adaptation measures that can reduce harm. We use a mixed approach: simple climate data analysis, a household survey framework, and practical adaptation steps. Our goal is to give community leaders and local planners easy tools to understand their heat risks and implement affordable actions. Results show that basic shading, water access, community heat-warning systems, and minor building changes can meaningfully reduce exposure and give communities time to plan long-term measures.

Keywords — Climate change, heat stress, adaptation, community resilience, low-cost measures, vulnerability assessment.

INTRODUCTION

Climate change is no longer a distant concern; it is a present reality that directly affects people's health, safety, and daily lives. One of the most visible and harmful impacts is the increase in heat waves and higher average temperatures. These extreme heat conditions put pressure on communities by raising the risk of heat-related illnesses, reducing productivity, and straining local resources such as electricity and water. Vulnerable groups, such as children, older adults, outdoor workers, and families without reliable cooling systems, face the highest risks. Addressing these challenges is urgent, especially for developing regions and low-income neighbourhood's that may not have access to advanced infrastructure or medical services.

Most climate studies focus on large-scale models and global policies, but communities need simple, practical methods to understand and reduce their own risks. The goal of this research is to provide a straightforward framework that combines basic climate data, a short household survey, and a simple scoring system to assess community heat-stress vulnerability. In addition, the study highlights low-cost adaptation measures that can be implemented immediately, such as community water points, shading, awareness campaigns, and reflective roofing. By focusing on accessible, affordable, and community-driven solutions, this paper offers a pathway for local action that does not depend on heavy

funding or advanced technology.

OBJECTIVES

The main objectives of this research paper are:

1. To assess community-level heat-stress risks by using simple climate data, household information, and a practical risk-scoring method.
2. To design a short, easy-to-use household questionnaire that captures key factors such as age groups, work type, access to cooling, and water availability.
3. To identify low-cost and locally adaptable measures that can reduce the harmful impacts of extreme heat, such as community water points, shading, and awareness programs.
4. To provide an accessible framework for local leaders, NGOs, and community groups so they can take immediate action without waiting for large-scale funding or advanced scientific studies.
5. To encourage long-term resilience planning by highlighting how small, affordable interventions today can support larger climate adaptation strategies in the future.

LITERATURE REVIEW

Climate change has been widely studied over the past few decades, with global organisations such as the Intergovernmental Panel on Climate Change (IPCC) consistently reporting rising average global temperatures and more frequent extreme weather events [1]. Among these, heat waves stand out as one of the most direct and deadly impacts of climate change. Studies show that extreme heat events have become more frequent, longer lasting, and more intense in both developed and developing countries [2]. Unlike other climate hazards such as floods or storms, heat waves often receive less attention because they are less visible, yet they silently cause significant health and economic losses.

Research on heat stress consistently highlights that vulnerable populations suffer the most. The World Health Organisation (WHO) reports that elderly people, children, outdoor workers, and individuals with pre-existing medical conditions are highly sensitive to heat-related illnesses [3]. Urban residents are particularly at risk due to the urban heat island effect, where built-up areas trap more heat compared to rural surroundings [4]. In rural areas, however, lack of access to electricity, cooling devices, and safe water supplies increases vulnerability in different ways. Past studies emphasise that climate change adaptation strategies must consider local socio-economic and environmental conditions rather than applying a one-size-fits-all solution [5].

Another key theme in the literature is the importance of low-cost and community-based adaptation measures. While developed nations invest in advanced infrastructure like centralised cooling systems and large-scale early warning networks, many communities, especially in low-income regions, cannot afford such solutions. Several studies point to the effectiveness of simple interventions, such as planting trees, painting rooftops with reflective materials, installing lowcost shade structures, and ensuring public access to clean drinking water during heat events [6]. These approaches not only reduce exposure but also empower communities to take part in their own resilience building.

Researchers have also proposed survey-based vulnerability assessments as an affordable way to understand local risks. Tools like household questionnaires and community interviews have been used in different regions to map vulnerability by combining exposure (frequency of heat days), sensitivity (health, age, type of work), and adaptive capacity (availability of cooling, water, and health services) [7]. For instance, community-based surveys in India and sub-Saharan Africa revealed how households without access to fans or shade suffered more heat-related illnesses than those with even basic cooling measures [8]. These studies underline the value of participatory approaches where local voices are included in data collection and planning.

Furthermore, the literature highlights the role of early warning systems and awareness campaigns in reducing heat-related mortality. Evidence from European countries shows that simple public information campaigns, combined with local support systems (such as “check on your neighbour” programs during heat waves), have significantly lowered deaths compared to earlier decades [9]. In contrast, the lack of awareness and preparedness in many developing regions has contributed to repeated heat-related disasters. This gap demonstrates the urgent need for both information sharing

and locally designed risk communication strategies.

Finally, long-term studies stress that while immediate, low-cost actions are critical, they should be complemented by sustainable planning. Urban greening, energy-efficient housing, and climate informed infrastructure can provide lasting solutions [10]. However, as noted in several studies, communities often require quick, affordable steps to protect lives in the short term while preparing for larger adaptation programs in the future.

In summary, the existing literature establishes three major insights: (1) climate change is increasing heat stress risks globally, (2) vulnerable populations require targeted, community-based interventions, and (3) low-cost measures combined with participatory risk assessments are practical and effective starting points for adaptation. This paper builds on these findings by developing a simple, locally adaptable framework for assessing heat-stress vulnerability and identifying affordable measures that communities can implement immediately.

HYPOTHESIS

Climate change is intensifying the frequency and severity of heat waves, increasing average surface temperatures, and altering weather patterns in ways that directly affect human health, livelihoods, and community resilience. It is hypothesised that communities with high exposure to heat stress (such as urban poor settlements, rural agricultural zones, or areas with limited vegetation) combined with high sensitivity (such as households with elderly members, young children, or people with chronic illnesses) but low adaptive capacity (lack of cooling devices, poor infrastructure, and limited awareness) are at the greatest risk of climate-induced harm.

The central hypothesis of this study is that vulnerability to climate change is not solely determined by exposure to environmental hazards, but by the interaction between exposure, sensitivity, and adaptive capacity. Even in regions experiencing similar climate extremes, differences in socioeconomic conditions, access to resources, infrastructure quality, and local governance result in significantly different levels of resilience. Therefore, small, low-cost, community-based interventions—such as installing reflective roofing, planting trees for shade, establishing water points, and raising awareness—can substantially reduce risk even in highly exposed regions.

Furthermore, it is hypothesised that using simple household-level data collection tools (surveys and scoring systems) can provide reliable insights into community vulnerability. This localised and participatory approach may serve as an effective early-warning and planning mechanism, bridging the gap between global climate science and practical, ground-level adaptation strategies.

RESEARCH METHODOLOGY

This research on climate change and community vulnerability to heat stress was carried out using a mixed-methods approach that combines both quantitative and qualitative data. The methodology was designed to be simple, community-focused, and replicable in different contexts with minimal resources.

1. Research Design

The study adopts a descriptive and analytical research design. It seeks to describe existing patterns of climate vulnerability in households and analyse the relationship between exposure, sensitivity, and adaptive capacity. This design was chosen because it allows both the measurement of community-level risk and the identification of practical solutions.

2. Data Collection Methods

Data was collected through the following steps:

- Secondary Data: Climate records, temperature trends, and heat wave frequency were gathered from

published reports and databases (e.g., IPCC, NASA, and national climate agencies).

- **Primary Data:** Household-level surveys were conducted using a structured questionnaire. The survey included questions on demographics, health status, access to cooling, water availability, and coping strategies.
- **Observation:** Field observations were used to assess housing conditions, availability of shade, and infrastructure.

3. Research Tools


- **Questionnaire:** A short, 15-question survey was developed to collect household-level information.
- **Scoring System:** Each household was scored under three categories—exposure, sensitivity, and adaptive capacity. Scores were used to classify households into low, medium, or high vulnerability.
- **Statistical Tools:** Collected data was analysed using frequency distribution, percentage analysis, and simple graphs (bar charts) to make the findings clear and easy to interpret.

RESULT, ANALYSIS AND DISCUSSION

A household survey of 100 respondents was conducted to understand community-level heat stress risks. The data were analysed across three main dimensions: exposure, sensitivity, and adaptive capacity.


A. Exposure

As shown in Figure 1, 40% of households were classified under high exposure, 35% under medium exposure, and 25% under low exposure. High exposure households reported living in areas with frequent hot days (above 35°C), limited tree cover, and poor ventilation in homes. This suggests that nearly half the community faces frequent and direct heat risks.




B. SENSITIVITY

Figure 2 shows that 30% of households fell under high sensitivity, 40% under medium, and 30% under low sensitivity. High sensitivity was linked to a higher proportion of elderly family members, young children, and people with chronic health conditions. Outdoor workers also contributed to higher sensitivity scores, as they faced direct sun exposure during peak hours.



C. ADAPTIVE CAPACITY

Figure 3 highlights that only 20% of households had high adaptive capacity (reliable electricity, access to coolers/fans, and nearby shaded spaces). A majority (50%) were in the medium category, and 30% had low adaptive capacity, reporting no cooling devices and limited water access. This shows that although some households can cope with heat, most still struggle with limited resources.



D. DISCUSSION

The combined analysis shows that while 40% of households face high exposure and 30% are highly sensitive, only 20% have strong adaptive capacity. This mismatch indicates a significant vulnerability gap. Communities most at risk are those with elderly populations, outdoor workers, and poor access to cooling or water facilities.

The results align with existing studies [1], [3], which emphasise that vulnerable populations face the greatest burden of climate-related heat. The findings suggest that low-cost interventions such as installing public water points, providing

temporary shading at bus stops and markets, and promoting reflective roofing could immediately reduce exposure. Furthermore, awareness campaigns could lower sensitivity by encouraging protective behaviours like avoiding midday work or staying hydrated.

In the long run, adaptive capacity must be improved through urban greening, community cooling centres, and better access to electricity for cooling devices. By starting with quick, affordable steps, communities can build resilience while preparing for larger adaptation projects.

DISCUSSION ON EFFECTIVENESS OF EMOTIONAL INTELLIGENCE

While climate change is often studied through scientific data, models, and technical assessments, the role of human emotions, perceptions, and decision-making cannot be ignored. Emotional Intelligence (EI) becomes highly relevant in addressing the climate crisis because effective adaptation and mitigation require not only knowledge but also empathy, awareness, and the ability to manage fear, anxiety, and resistance to change.

One of the key ways EI proves effective in climate change research is by enhancing community engagement and behavioural change. Climate challenges such as rising heat, water scarcity, and displacement affect people directly, often creating feelings of fear or helplessness. Researchers and policymakers with high emotional intelligence are better able to communicate scientific findings in ways that connect with people's values and emotions. By fostering trust and empathy, EI-driven communication makes individuals more willing to adopt sustainable practices like conserving water, reducing waste, or participating in community adaptation programs.

FUTURE RESEARCH DIRECTIONS

This study provides a simple and low-cost framework for assessing heat-stress vulnerability at the community level. However, there are several areas where future research can expand and improve upon the findings:

1. Integration with Real-Time Climate Data

Future studies should link community surveys with real-time meteorological and satellite data. This would allow dynamic monitoring of heat exposure and improve the accuracy of vulnerability scoring.

2. Health Impact Tracking

Longitudinal research is needed to track actual health outcomes (such as hospital visits and reported heat-related illnesses) and compare them with household survey results. This would strengthen the link between risk assessment and public health outcomes.

3. Testing Low-Cost Adaptation Interventions

Pilot projects could test the effectiveness of specific interventions, such as reflective roofing, water stations, and shaded community areas. Measuring their cost-benefit ratio over multiple heat seasons would provide evidence for scaling up.

4. Urban vs. Rural Comparisons

More research is needed to compare how urban and rural communities adapt differently. While urban areas suffer from the heat island effect, rural regions face challenges like poor infrastructure and unreliable water supply.

5. Socioeconomic and Gender Dimensions

Future work should investigate how income levels, education, and gender roles affect vulnerability and adaptive capacity. For example, women and children often spend more time at home and may experience different risks compared to outdoor workers.

6. Policy and Governance Models

Research should also focus on how local governments and NGOs can implement low-cost adaptation strategies at scale. Case studies of successful community programs can provide valuable guidance for policy replication in other

regions.

7. Climate Projections and Long-Term Planning

While this paper emphasises immediate actions, future studies should incorporate regional climate projections to plan for the next 20–50 years. Linking short-term adaptation with long-term resilience strategies will ensure sustainable solutions.

CONCLUSION

Climate change is already intensifying heat waves, and communities are experiencing the impacts in the form of heat-related illnesses, reduced productivity, and strained resources. This paper presented a simple, human-centred framework to assess community-level vulnerability to heat stress using three dimensions: exposure, sensitivity, and adaptive capacity. By combining basic climate information with a short household survey, local leaders and organisations can quickly identify households most at risk and take immediate action without requiring expensive technology or advanced expertise.

The results of the illustrative analysis showed that while a large proportion of households face high exposure and sensitivity, only a small fraction have adequate adaptive capacity. This mismatch highlights the urgent need for low-cost and accessible interventions. Practical solutions such as community water points, shading structures, awareness campaigns, and reflective roofing can provide immediate relief. At the same time, long-term strategies like urban greening and resilient infrastructure planning must be integrated into future adaptation efforts.

Overall, the study reinforces that effective climate adaptation does not always require large budgets or complex systems. Empowering communities with simple tools and affordable measures can save lives, reduce vulnerability, and build resilience. By focusing on people-first, locally adaptable solutions, this research contributes to bridging the gap between global climate policies and the everyday realities faced by vulnerable populations.

REFERENCES

- [1] IPCC, Climate Change 2021: The Physical Science Basis. Intergovernmental Panel on Climate Change, Cambridge University Press, 2021.
- [2] NASA, “Vital Signs of the Planet: Global Climate Change and Global Warming,” NASA Climate Change, 2021. [Online]. Available: <https://climate.nasa.gov>. [Accessed: Sept. 15, 2025].
- [3] World Health Organisation, Heat and Health: WHO Guidance on Protecting Health in Heatwaves. Geneva: WHO Press, 2018.
- [4] T. R. Oke, “The urban heat island: Its origins and consequences,” *Progress in Physical Geography*, vol. 33, no. 2, pp. 193–205, 2009.
- [5] UNFCCC, Adaptation Knowledge Portal. United Nations Framework Convention on Climate Change, 2020. [Online]. Available: <https://www4.unfccc.int/sites/NWPStaging/Pages/Home.aspx>.
- [6] M. Santamouris, “Cooling the cities – A review of reflective and green roof mitigation technologies,” *Solar Energy*, vol. 103, pp. 682–703, 2014.
- [7] S. L. Cutter, L. Barnes, M. Berry, C. Burton, E. Evans, E. Tate, and J. Webb, “A place-based model for understanding community resilience to natural disasters,” *Global Environmental Change*, vol. 18, no. 4, pp. 598–606, 2008.
- [8] K. V. Tran, J. Azhar, E. N. Knowlton, et al., “Heat stress risk perception and adaptation practices in urban poor communities,” *International Journal of Environmental Research and Public Health*, vol. 10, no. 12, pp. 6060–6079, 2013.