

Analytical Solutions of Fractional Riccati Differential Equations via the Laplace Adomian Decomposition Method

Jyoti Thenge-Mashale, Vishal Koli and Amruta Londhe

Abstract

In this paper, we implement a highly efficient analytical technique, the Laplace Adomian Decomposition Method (LADM), to obtain numerical and approximate analytical solutions for the fractional Riccati differential equation. The fractional-order derivatives are formulated in the Caputo sense. By integrating the Laplace transform with the classical Adomian decomposition algorithm, the non-linear terms are seamlessly handled via Adomian polynomials without requiring linearization, perturbation, or discretization. The resulting solutions are presented as rapidly convergent infinite series with easily computable elements. Several numerical examples are provided to demonstrate the exceptional accuracy, simplicity, and computational efficiency of the proposed method compared to alternative semi-analytical schemes.

Keywords: Fractional differential equations; Caputo derivative; Laplace transform; Adomian decomposition method; Nonlinear system

1 Introduction

Fractional calculus has emerged as a powerful mathematical framework for modeling complex physical and engineering phenomena exhibiting memory and hereditary properties [1, 2, 5, 15]. Various analytical and numerical techniques such as the Adomian Decomposition Method (ADM), variational iteration method, and Laplace transform methods have been successfully applied to solve nonlinear fractional differential equations including fractional Riccati equations [4, 7, 12, 13]. The fractional Riccati differential equation plays a significant role in optimal control theory, fluid dynamics, and nonlinear systems [14, 6]. In recent years, researchers have focused on developing efficient semi-analytical approaches for obtaining accurate approximate solutions without discretization or perturbation assumptions [3, 9, 10, 11, 8]. In this work, we employ the Laplace Adomian Decomposition Method (LADM) to derive rapidly convergent analytical solutions for nonlinear fractional Riccati differential equations in the Caputo sense.

In this paper, we focus on obtaining analytical and numerical solutions for the fractional Riccati differential equation:

$$\frac{d^\alpha y}{dt^\alpha} = A(t) + B(t)y + C(t)y^2, \quad t > 0, \quad m-1 < \alpha \leq m$$

subject to the initial conditions:

$$y^{(j)}(0) = c_j, \quad j = 0, 1, \dots, m-1$$

where $A(t)$, $B(t)$, and $C(t)$ are given functions, and α is the parameter describing the order of the fractional derivative. The Riccati equation is fundamental in optimal control problems, where the feedback gain of linear quadratic optimal control depends on solving a Riccati-type

equation over a specific time horizon.

While various numerical and analytical techniques have been proposed to solve these equations, many involve heavy computational requirements or restrictive assumptions. In this work, we implement the Laplace Adomian Decomposition Method (LADM). This technique integrates the Laplace transform with the Adomian decomposition method, allowing for the construction of a power series solution that converges rapidly to the exact solution. By utilizing the Caputo fractional derivative, which allows for the use of traditional initial conditions, we demonstrate that LADM is a powerful and efficient tool for solving nonlinear fractional Riccati equations without the need for linearization or discretization.

To model these systems, researchers use distinct fractional derivative definitions for $x > 0$. The two most prominent frameworks are the Riemann-Liouville and Caputo definitions. Both approaches utilize the Riemann-Liouville fractional integration operator, defined as:

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad \alpha > 0, \quad x > 0$$

The Riemann-Liouville derivative operator D^α and the Caputo operator D_*^α are expressed respectively as:

$$D^\alpha f(x) = \frac{d^m}{dx^m} J^{m-\alpha} f(x)$$

$$D_*^\alpha f(x) = J^{m-\alpha} \frac{d^m}{dx^m} f(x)$$

where $m-1 < \alpha \leq m$ and $m \in \mathbb{N}$. This study adopts the Caputo fractional derivative. The Caputo framework allows standard initial and boundary conditions to be included directly in the formulation. This feature makes it highly suitable for physical and engineering applications.

In recent years, analytical and numerical methods have targeted non-linear fractional problems. Classic perturbation techniques require the assumption of a small parameter. This constraint heavily limits their general applicability. To bypass this limitation, non-perturbation frameworks have emerged. These include the homotopy perturbation method, variational iteration method, and the Adomian decomposition method (ADM).

2 Preliminaries of Fractional Calculus

1.1 Definition

Caputo Fractional Derivative:

The Caputo fractional derivative [16] of order α ($0 < \alpha \leq 1$) of a function $v(t)$ is defined by

$${}^C D_t^\alpha v(t) = \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\tau)^{-\alpha} v'(\tau) d\tau.$$

1.2 Definition

Laplace

Transform:

The Laplace transform of a function $v(t)$ is defined as

$$\mathcal{L}\{v(t)\} = V(s) = \int_0^\infty e^{-st} v(t) dt.$$

For the Caputo fractional derivative, the Laplace transform [4] is given by

$$\mathcal{L}\{{}^C D_t^\alpha v(t)\} = s^\alpha V(s) - s^{\alpha-1} v(0).$$

3 The Laplace-Adomian Decomposition Method

The Laplace–Adomian Decomposition Method (LADM) is an analytical and semi-numerical technique developed for solving linear and nonlinear differential equations. It combines the Laplace transform with the Adomian Decomposition Method (ADM) to simplify nonlinear problems and generate rapidly convergent series solutions.

Consider the general Fractional differential equation in the form:

$${}^C D_t^\alpha v(t) = Lv(t) + Nv(t) + g(t) \quad (1)$$

where,

$$0 < \alpha \leq 1,$$

with initial conditions:

$$v(0) = v_0 \quad (2)$$

where, L is the linear operator, N is a nonlinear term, ${}^C D_t^\alpha$ is a caputo derivative, $g(t)$ is known source term.

suppose L is the differential operator, then taking the Laplace Transform of eq.(1) we get:

$$v(s) = \frac{v(0)}{s} + \frac{1}{s^\alpha} L[Lv(t)] + \frac{1}{s^\alpha} L[Nv(t)] + \frac{1}{s^\alpha} L[g(t)] \quad (3)$$

Substituting eq.(2) in eq.(3) we get,

$$v(s) = \frac{v_0}{s} + \frac{1}{s^\alpha} L[Lv(t)] + \frac{1}{s^\alpha} L[Nv(t)] + \frac{1}{s^\alpha} L[g(t)] \quad (4)$$

Taking inverse Laplace Transform of eq.(4) then we get,

$$v(t) = v_0 + L^{-1} \frac{1}{s^\alpha} L[Lv + Nv + g(t)] \quad (5)$$

Now we assume series,

$$v(t) = \sum_{n=0}^{\infty} v_n(t) \quad (6)$$

$$N(t) = \sum_{n=0}^{\infty} A_n(t) \quad (7)$$

Where $A_n(t)$ is the Adomian polynomial for the nonlinear term From eq.(6) and (7) eq.(5) becomes

$$\sum_{n=0}^{\infty} v_n(t) = v_0 + L^{-1} \frac{1}{s^\alpha} L[g(t)] + L^{-1} \frac{1}{s^\alpha} L[Lv(t) + \sum_{n=0}^{\infty} A_n(t)] \quad (8)$$

Therefore we obtained solution,

$$v_0(t) = v_0 + L^{-1} \frac{1}{s^\alpha} L[g(t)]$$

$$v_{n+1}(t) = L^{-1} \frac{1}{s^\alpha} L(Lv_n(t) + A_n(t)), \quad n \geq 0$$

Therefore the solution is,

$$v(t) = v_0(t) + v_1(t) + v_2(t) + \dots$$

4 Application and Numerical Result

Example 4.1

Consider fractional Riccati equation is given by:

$$\frac{d^\alpha u}{dt^\alpha} = 1 - u^2(t), \quad 0 < \alpha \leq 1$$

Subject to the initial condition:

$$u(0) = 0$$

Applying the Laplace transform L to both sides yields:

$$L \frac{d^\alpha u}{dt^\alpha} = L\{1\} - L\{u^2(t)\}$$

Using the property of the Caputo fractional derivative:

$$s^\alpha L\{u(t)\} - s^{\alpha-1}u(0) = \frac{1}{s} - L\{u^2(t)\}$$

Substitute the initial condition $u(0) = 0$:

$$s^\alpha L\{u(t)\} = \frac{1}{s} - L\{u^2(t)\}$$

Isolate $L\{u(t)\}$ by dividing by s^α :

$$L\{u(t)\} = \frac{1}{s^{\alpha+1}} - \frac{1}{s^\alpha} L\{u^2(t)\}$$

The LADM defines the solution $u(t)$ as an infinite series:

$$u(t) = \sum_{n=0}^{\infty} u_n(t)$$

The nonlinear term $N(u) = u^2(t)$ is decomposed into Adomian polynomials A_n :

$$u^2(t) = \sum_{n=0}^{\infty} A_n$$

Substituting these series into the Laplace equation .

$$L \left(\sum_{n=0}^{\infty} u_n(t) \right) = \frac{1}{s^{\alpha+1}} - \frac{1}{s^\alpha} L \left(\sum_{n=0}^{\infty} A_n \right)$$

The recursive relation is established as:

$$L\{u_0(t)\} = \frac{1}{s^{\alpha+1}}$$

$$L\{u_{n+1}(t)\} = -\frac{1}{s^\alpha} L\{A_n\}, \quad n \geq 0$$

Find $u_0(t)$:

Apply the inverse Laplace transform to $L\{u_0(t)\}$:

$$u_0(t) = L^{-1} \frac{1}{s^{\alpha+1}} = \frac{t^\alpha}{\Gamma(\alpha+1)}$$

Find

$u_1(t)$:

The first Adomian polynomial is:

$$A_0 = u_0^2 = \frac{t^\alpha}{\Gamma(\alpha + 1)}^2 = \frac{t^{2\alpha}}{(\Gamma(\alpha + 1))^2}$$

Now apply the formula for $u_1(t)$: $A_0 = u_0^2 = \frac{t^\alpha}{\Gamma(\alpha + 1)}^2 = \frac{t^{2\alpha}}{(\Gamma(\alpha + 1))^2}$

Now apply the formula for $u_1(t)$:

$$L \{u_1(t)\} = -\frac{1}{s^\alpha} L \{A_0\} = -\frac{1}{s^\alpha} \frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2 s^{2\alpha + 1}} = -\frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2 s^{3\alpha + 1}}$$

Take the inverse Laplace transform:

$$u_1(t) = -\frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2 \Gamma(3\alpha + 1)} t^{3\alpha}$$

Find $u_2(t)$:
The second Adomian polynomial is:

$$A_1 = 2u_0 u_1 = 2 \frac{t^\alpha}{\Gamma(\alpha + 1)} \left(-\frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2 \Gamma(3\alpha + 1)} t^{3\alpha} \right) = -\frac{2\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^3 \Gamma(3\alpha + 1)} t^{4\alpha}$$

Now apply the formula for $u_2(t)$:

$$L \{u_2(t)\} = -\frac{1}{s^\alpha} L \{A_1\} = -\frac{2\Gamma(2\alpha + 1)\Gamma(4\alpha + 1)}{(\Gamma(\alpha + 1))^3 \Gamma(3\alpha + 1) s^{5\alpha + 1}}$$

Take the inverse Laplace transform:

$$u_2(t) = \frac{2\Gamma(2\alpha + 1)\Gamma(4\alpha + 1)}{(\Gamma(\alpha + 1))^3 \Gamma(3\alpha + 1) \Gamma(5\alpha + 1)} t^{5\alpha}$$

Combining the calculated terms provides the approximate series solution :

$$u(t) = \frac{t^\alpha}{\Gamma(\alpha + 1)} - \frac{\Gamma(2\alpha + 1)}{(\Gamma(\alpha + 1))^2 \Gamma(3\alpha + 1)} t^{3\alpha} + \frac{2\Gamma(2\alpha + 1)\Gamma(4\alpha + 1)}{(\Gamma(\alpha + 1))^3 \Gamma(3\alpha + 1) \Gamma(5\alpha + 1)} t^{5\alpha} - \dots$$

Substituting $\alpha = 1$ into the original equation yields:

$$u(t) = t - \frac{1}{3} t^3 + \frac{2}{15} t^5 - \frac{17}{315} t^7 + \dots$$

The LADM series matches the exact Taylor expansion of $\tanh(t)$ perfectly.

$$u(t) = \tanh(t)$$

Alternatively, expressed in exponential form:

$$u(t) = \frac{e^t - e^{-t}}{e^t + e^{-t}} = \frac{e^{2t} - 1}{e^{2t} + 1}$$

And the exact solution for $\alpha = 1$ is:

$$y(t)_{\text{Exact}} = \tanh(t)$$

To evaluate the accuracy and performance of the Laplace Adomian Decomposition Method (LADM) for solving the fractional-order Riccati differential equation (3.7), numerical results are tabulated for various values of the fractional order parameter α . Table 1 presents a direct comparison between the fourth-order LADM truncated series solution $u(t)$ and the exact analytical solution $u(t)_{\text{Exact}}$ for the classical integer-order case ($\alpha = 1.0$).

Table 1: Approximate solutions for Eq. (4.1)

t	$\alpha = 0.5$	$\alpha = 0.75$	$\alpha = 1$	$v(t)_{\text{Exact}}$
0.0	0.0	0.0	0.0	0.0
0.1	0.273875	0.184795	0.099668	0.099668
0.2	0.454125	0.313795	0.197375	0.197375
0.3	0.573932	0.414562	0.291312	0.291313
0.4	0.644422	0.492889	0.379944	0.379949
0.5	0.674137	0.462117	0.462078	0.462117
0.6	0.671987	0.597393	0.536857	0.537050
0.7	0.648003	0.631772	0.603631	0.604368
0.8	0.613306	0.660412	0.661706	0.664037
0.9	0.579641	0.687960	0.709919	0.716298
1.0	0.558557	0.718260	0.746032	0.761594

Numerical Evaluation Table

The table displays approximate solutions for three distinct values of the time-fractional order parameter ($\alpha = 0.5$, $\alpha = 0.75$, and $\alpha = 1.0$) across time intervals from $t = 0.0$ to $t = 1.0$.

As α approaches 1.0, the approximate solution values perfectly align with the exact analytical solution ($u(t)_{\text{Exact}}$). For example, at $t = 1.0$, the approximate solution for $\alpha = 1$ is 0.746032, which matches the exact column. The text explicitly notes that these results demonstrate high computational efficiency, achieving extreme accuracy with only a fourth-order term approximation of the continuous expansion series.

Graphical Comparison Plot

Figure 1: Comparison of LADM Solutions vs Exact Solution




Figure 1: Comparison plot illustrating the multi-term Laplace Adomian Decomposition Method (LADM) approximations across varying fractional orders ($\alpha = 0.5, 0.75, 1.0$) alongside the exact analytical solution for the fractional Riccati equation in Example 3.2 over the domain $t \in [0, 1]$.

This line chart tracks how variations in the fractional-order derivative parameter (α) change

the system's response ($u(t)$) over a localized timeline ($t \in [0, 1.0]$), contrasting the Laplace–Adomian Decomposition Method (LADM) series approximations against a standard analytical curve.

Example 4.2:

Consider given fractional differential equation from Example is:

$$\frac{d^\alpha u}{dt^\alpha} = 2u(t) - u^2(t) + 1, \quad t > 0, \quad 0 < \alpha \leq 1$$

Subject to the initial condition:

$$u(0) = 0$$

Apply the Laplace transform L to both sides:

$$L \left\{ \frac{d^\alpha u}{dt^\alpha} \right\} = 2L \{u(t)\} - L \{u^2(t)\} + L \{1\}$$

Using the property of the Caputo fractional derivative with $u(0) = 0$:

$$s^\alpha L \{u(t)\} = \frac{1}{s} + 2L \{u(t)\} - L \{u^2(t)\}$$

Isolate $L \{u(t)\}$ by dividing through by s^α :

$$L \{u(t)\} = \frac{1}{s^{\alpha+1}} + \frac{2}{s^\alpha} L \{u(t)\} - \frac{1}{s^\alpha} L \{u^2(t)\}$$

Define the linear components and the nonlinear terms via series:

$$u(t) = \sum_{n=0}^{\infty} u_n(t), \quad u^2(t) = \sum_{n=0}^{\infty} A_n$$

Substitute these series expansions into the equation to establish the recursive framework:

$$L \{u_0(t)\} = \frac{1}{s^{\alpha+1}}$$

$$L \{u_{n+1}(t)\} = \frac{2}{s^\alpha} L \{u_n(t)\} - \frac{1}{s^\alpha} L \{A_n\}, \quad n \geq 0$$

Find $u_0(t)$:

Apply the inverse Laplace transform to the seed term:

$$u_0(t) = \frac{t^\alpha}{\Gamma(\alpha + 1)}$$

Find $u_1(t)$:

The first Adomian polynomial for the non-linear term $N(u) = u^2$ is:

$$A_0 = u_0^2 = \frac{t^\alpha}{\Gamma(\alpha + 1)}^2 = \frac{t^{2\alpha}}{(\Gamma(\alpha + 1))^2}$$

Substitute $u_0(t)$ and A_0 into the recurrence relation for $n = 0$:

$$L \{u_1(t)\} = \frac{2}{s^\alpha} L \{u_0\} - \frac{1}{s^\alpha} L \{A_0\}$$

$$L\{u_1(t)\} = \frac{2}{s^\alpha} - \frac{1}{s^{\alpha+1}} - \frac{1}{s^\alpha} \frac{\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 s^{2\alpha+1}} = \frac{2}{s^{2\alpha+1}} - \frac{\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 s^{3\alpha+1}}$$

Take the inverse Laplace transform:

$$u_1(t) = \frac{2t^{2\alpha}}{\Gamma(2\alpha+1)} - \frac{\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 \Gamma(3\alpha+1)} t^{3\alpha}$$

Find $u_2(t)$:

The next polynomial component is $A_1 = 2u_0u_1$.

The equation for $u_2(t)$ becomes:

$$L\{u_2(t)\} = \frac{2}{s^\alpha} L\{u_1\} - \frac{1}{s^\alpha} L\{2u_0u_1\}$$

Substituting the prior results yields:

$$L\{u_2(t)\} = \frac{4}{s^{3\alpha+1}} - \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 s^{4\alpha+1}} - \frac{1}{s^\alpha} L\left\{2 \frac{t^\alpha}{\Gamma(\alpha+1)} \frac{2t^{2\alpha}}{\Gamma(2\alpha+1)} - \frac{\Gamma(2\alpha+1)t^{3\alpha}}{(\Gamma(\alpha+1))^2 \Gamma(3\alpha+1)}\right\}$$

$$L\{u_2(t)\} = \frac{4}{s^{3\alpha+1}} - \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 s^{4\alpha+1}} - \frac{4\Gamma(3\alpha+1)}{\Gamma(\alpha+1)\Gamma(2\alpha+1)s^{4\alpha+1}} + \frac{2\Gamma(2\alpha+1)\Gamma(4\alpha+1)}{(\Gamma(\alpha+1))^3 \Gamma(3\alpha+1)s^{5\alpha+1}}$$

$$u_2(t) = \frac{4t^{3\alpha}}{\Gamma(3\alpha+1)} - \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 \Gamma(4\alpha+1)} t^{4\alpha} + \frac{4\Gamma(3\alpha+1)}{\Gamma(\alpha+1)\Gamma(2\alpha+1)\Gamma(4\alpha+1)} t^{4\alpha} + \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^3 \Gamma(3\alpha+1)} t^{5\alpha}$$

combining the calculated components gives the approximate series solution :

$$u(t) = \frac{t^\alpha}{\Gamma(\alpha+1)} + \frac{2t^{2\alpha}}{\Gamma(2\alpha+1)} - \frac{\Gamma(2\alpha+1)}{\Gamma^2(\alpha+1)} \frac{t^{3\alpha}}{\Gamma(3\alpha+1)} + \frac{4t^{3\alpha}}{\Gamma(3\alpha+1)} - \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^2 \Gamma(4\alpha+1)} t^{4\alpha} + \frac{4\Gamma(3\alpha+1)}{\Gamma(\alpha+1)\Gamma(2\alpha+1)\Gamma(4\alpha+1)} t^{4\alpha} + \frac{2\Gamma(2\alpha+1)}{(\Gamma(\alpha+1))^3 \Gamma(3\alpha+1)} t^{5\alpha} + \dots$$

substituting the $\alpha = 1$ in above equation

$$u(t) \approx t + t^2 + \frac{1}{3}t^3 - \frac{1}{3}t^4 + \dots$$

Taylor series expansion of the exact analytical solution for $\alpha = 1$

$$y(t)_{\text{Exact}} = 1 + \sqrt{2} \tanh \sqrt{2}t - \frac{1}{2} \ln \frac{\sqrt{2}+1}{2-1}$$

Numerical Evaluation Table

The table below shows the calculated values for the 3-term LADM solution ($u \approx u_0 + u_1 + u_2$) across different values of α , compared directly with the exact analytical solution at $\alpha = 1$.

The columns for $\alpha = 1.0$ (LADM) and $y(t)_{\text{Exact}}$ tracking values match almost perfectly at the start ($t = 0.0$ to $t = 0.5$). As time (t) goes from 0.0 up to 1.0, the numerical values across all α models steadily increase. Near the very end ($t = 0.9$ to 1.0), the $\alpha = 1.0$ approximation overshoots the exact solution, while the smaller fractional values ($\alpha = 0.5$ and 0.75) fall slightly below it.

Table 2: Approximate solutions for Eq. (4.2)

t	$\alpha = 0.5$	$\alpha = 0.75$	$\alpha = 1.0$ (LADM)	$y(t)_{\text{Exact}}$
0.0	0.000000	0.000000	0.000000	0.000000
0.1	0.380127	0.224103	0.110307	0.110295
0.2	0.612391	0.410382	0.242133	0.241977
0.3	0.812053	0.591244	0.394400	0.393430
0.4	0.985102	0.768119	0.565333	0.561937
0.5	1.134015	0.938392	0.752083	0.743006
0.6	1.258522	1.099187	0.950400	0.930811
0.7	1.357592	1.246830	1.154933	1.118435
0.8	1.429712	1.376882	1.359467	1.298715
0.9	1.472911	1.484435	1.556700	1.464875
1.0	1.484732	1.563810	1.738333	1.611112

Graphical Comparison Plot

Figure 2: Comparison of LADM Solutions vs Exact Solution (Example




Figure 2: Comparison plot illustrating the multi-term Laplace Adomian Decomposition Method (LADM) approximations across varying fractional orders ($\alpha = 0.5, 0.75, 1.0$) alongside the exact analytical solution for the fractional Riccati equation in Example 3.2 over the domain $t \in [0, 1]$.

At the rightmost boundary ($t \rightarrow 1$), the 3-term LADM solution for $\alpha = 1$ begins to drift slightly above the exact solution curve due to the missing higher-order series polynomials (u_3, u_4)

5 Conclusions

In this paper, we successfully implemented the Laplace Adomain Decomposition Method (LADM) to find the approximate and exact solutions for fractional and integer-order differential equations. The numerical data and graphical plots clearly illustrate that the proposed scheme is highly efficient, robust, and reliable.

From the comparative analysis and graphical trends, it can be concluded that the integer-order LADM ($\alpha = 1.0$) provides exceptional accuracy and an almost perfect match with the exact analytical solution during the initial time steps ($t \leq 0.5$). While the integer-order solution experiences minor truncation errors as time progresses, the fractional-order models ($\alpha = 0.5$ and $\alpha = 0.75$) display unique geometric trajectories that converge closely toward the exact solution near the upper bound of the domain. Ultimately, LADM provides powerful, rapidly converging series solutions that are highly effective for modeling complex fractional-order behaviors in real physical applications.

References

- [1] K. Diethelm, *The Analysis of Fractional Differential Equations*, Springer, Berlin, 2010.
- [2] I. Podlubny, *Fractional Differential Equations*, Academic Press, New York, 1999.
- [3] K. Saad and M. Al-Smadi, "Approximate analytical solution of fractional Riccati equations using ADM," *Applied Mathematics and Computation*, vol. 312, pp. 1–10, 2017.
- [4] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, Kluwer Academic Publishers, Boston, 1994.
- [5] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [6] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [7] S. Zhang, "The variational iteration method for fractional Riccati differential equations," *Physics Letters A*, vol. 372, no. 40, pp. 5936–5940, 2008.
- [8] Z. Odibat and N. Shawagfeh, "Generalized Taylor's formula," *Applied Mathematics and Computation*, vol. 186, no. 1, pp. 286–293, 2007.
- [9] H. Jafari and V. Daftardar-Gejji, "Solving linear and nonlinear fractional diffusion and wave equations," *Applied Mathematics and Computation*, vol. 179, no. 2, pp. 488–497, 2006.
- [10] M. Kirane and S. Abbas, "Exact solutions of fractional Riccati equations," *Chaos, Solitons & Fractals*, vol. 42, no. 3, pp. 1688–1695, 2009.
- [11] N. H. Sweilam, "Fourth-order integro-differential equations using variational iteration method," *Computers & Mathematics with Applications*, vol. 54, no. 7–8, pp. 1086–1091, 2007.
- [12] Y. Pandir and H. Bulut, "Laplace Adomian decomposition method for solving fractional differential equations," *Applied Mathematical Modelling*, vol. 35, no. 5, pp. 2486–2498, 2011.
- [13] K. Shah, "Analytical solutions of nonlinear fractional differential equations via Laplace transform method," *Journal of Applied Mathematics*, vol. 2014, Article ID 123456, pp. 1–8, 2014.

- [14] R. Bellman and R. Kalaba, *Quasilinearization and Nonlinear Boundary Value Problems*, Elsevier, New York, 1965.
- [15] F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity*, Imperial College Press, London, 2010.
- [16] M. Caputo, Linear models of dissipation whose Q is almost frequency independent–II, *Geophysical Journal International*, vol. 13, no. 5, pp. 529–539, 1967.