

Fungi as Environmental Guardians: Enzymatic Solutions to Pollution

Poonam¹, Kavita²

¹Assistant Professor, Guru Kashi University, Talwandi Sabo, District Bathinda, Punjab, 151302, India

²Assistant Professor, Guru Kashi University, Talwandi Sabo, District Bathinda, Punjab, 151302, India

Corresponding author: Poonam

Email id: poonamsain56229@gmail.com

ABSTRACT

Fungi grow simply, inexpensively and act like little factories. Their enzymes are safer and more environmentally friendly for everyone to use than dangerous chemicals. This review covers the use of these enzymes by scientists to remediate contaminated soil, treat wastewater and even convert waste to valuable products such as biofuels.

For example, fungi are great cleaners. They produce enzymes. These are tiny workers that decompose all sorts of nasty environmental stuff. It all boils down to those workhorses of fungal enzymes. We consider different types, e.g. starch amylases, wood ligninases and plant cellulases. These enzymes can decontaminate, such as pesticides from land and dyes from water.

This paper critically assesses the potential of fungal enzymes for wastewater treatment, the conversion of organic waste into bio-based products and environmental remediation.

However, there are considerable differences in the efficacy of these enzymes, thus optimisation is a step towards improving their catalytic abilities. Genetic engineering or optimisation of environmental parameters for enhanced fungal strains for increased enzyme production may contribute to improved enzyme kinetics. Finally, more research is needed to scale up and reduce the cost of using fungal enzymes and to make them more widely used to produce environmentally friendly remedies for environmental contamination. The key opportunity for advancing green technology for environmental remediation is to unlock the full potential of fungal enzymes.

INTRODUCTION

Pollution is placing huge stress on ecosystems, creating an urgent need for innovative and sustainable environmental management strategies. Natural biological systems, particularly fungi, offer promising alternatives to conventional remediation methods that often rely on large quantities of hazardous chemicals. Owing to their exceptional enzymatic capabilities, fungi can transform complex pollutants into simpler and less toxic compounds, making them effective biological catalysts for environmental cleanup. Fungal-based remediation, commonly known as mycoremediation, has emerged as an eco-friendly and cost-effective approach for restoring contaminated ecosystems and promoting environmental sustainability. (1-2)

Fungi secrete a wide range of extracellular enzymes, each specifically adapted to degrade different classes of pollutants. For instance, cellulases hydrolyze cellulose, a major component of plant biomass; ligninases break down lignin, a highly recalcitrant polymer present in woody materials; and amylases catalyze the degradation of starch. In addition, oxidative enzymes such as laccases, manganese peroxidases, and lignin peroxidases play crucial roles in the degradation of synthetic dyes, pharmaceuticals, pesticides, hydrocarbons, and other xenobiotic compounds (2-3).

Bioremediation has gained considerable attention as a green technology that employs living organisms to transform, detoxify, or remove contaminants from polluted environments. Among various biological agents, fungi are particularly promising because of their remarkable metabolic versatility, extensive mycelial networks, and ability to produce diverse extracellular and intracellular enzymes capable of degrading complex pollutants. Recent studies have highlighted the

importance of fungal consortia, plant–fungal interactions, and advanced omics technologies in enhancing pollutant degradation efficiency and field applicability(1).These enzymes work synergistically to form an effective enzymatic toolbox capable of degrading a broad spectrum of environmental contaminants, ranging from synthetic dyes in wastewater to persistent pesticides, pharmaceuticals, plastics, petroleum hydrocarbons, and heavy metals in soil and aquatic systems(1-4).

Fungal enzymes offer several advantages over traditional chemical treatment methods. They are biodegradable, renewable, environmentally benign, and generally function under mild operational conditions. Furthermore, the relatively low cost of fungal cultivation and the scalability of enzyme production make fungal biotechnology suitable for large-scale environmental applications. Recent advances in enzyme engineering, immobilization technologies, and bioreactor design have further improved the efficiency and practical feasibility of fungal-based remediation systems (3).

This article provides an overview of fungal enzymes and their applications in waste treatment and bioremediation, highlighting their growing importance as sustainable tools for combating environmental pollution and advancing green technologies.

Moreover, it shows the progress and prospects for the production of potent fungal enzymes and their broad use in the cleaning of polluted environments, paving the way for more sustainable and environmentally friendly environmental solutions.

TYPES OF FUNGAL ENZYMES USED IN ENVIRONMENTAL MANAGEMENT

Enzyme	Source Organism	Working Efficiency	Key Feature	Environmental Application	Properties	Additional Info	Reference no.
Laccase	<i>Trametes versicolor</i>	High	Oxidizes phenolic compounds	Bioremediation of wastewater	Optimal pH: 4-6, temperature stability: 40-60°C	Used in the breakdown of lignin	5,6
Lignin Peroxidase	<i>Phanerochaete chrysosporium</i>	High	Degrades lignin polymers	Wood pulp bleaching, bioremediation	Optimal pH: 3-5, temperature stability: 25-40°C	Breaks down complex lignin structures	7
Manganese Peroxidase	<i>Phanerochaete chrysosporium</i>	Moderate	Oxidizes lignin-related compounds	Bioremediation, waste treatment	Optimal pH: 3-5, temperature stability: 30-50°C	Used in the degradation of organic pollutants	8
Cellulase	<i>Trichoderma reesei</i>	Very High	Breaks down cellulose	Wastepaper recycling, soil management	Optimal pH: 4.5-5.5, temperature	Important for biodegradation of plant waste	9,10

					stability: 50-60°C		
Chitinase	<i>Colletotrichum graminicola</i>	Moderate	Degrades chitin (insects, fungi)	Agricultural pest control	Optimal pH: 5.5-7.5, temperature stability: 40-50°C	Involved in fungal cell wall degradation	11
Amylase	<i>Aspergillus niger</i>	High	Breaks down starches	Bioremediation, industrial use	Optimal pH: 4.5-5.5, temperature stability: 45-60°C	Used for detoxification of starch waste	12
Peroxidase	<i>Coriarius versicolor</i>	Moderate	Degrades organic compounds	Detoxification of phenols	Optimal pH: 5-6, temperature stability: 30-50°C	Effective in treating industrial effluents	13
Xylanase	<i>Thermomyces lanuginosus</i>	High	Breaks down xylans (hemicelluloses)	Biodegradation of plant waste	Optimal pH: 5-7, temperature stability: 50-80°C	Important for wood and paper industry	14
Lipase	<i>Candida rugosa</i>	Moderate	Breaks down lipids	Biodegradation of oils and fats	Optimal pH: 6-8, temperature stability: 35-50°C	Used in oil spill cleanup and waste treatment	15
Protease	<i>Aspergillus oryzae</i>	High	Breaks down proteins	Wastewater treatment	Optimal pH: 5.5-7, temperature stability: 45-55°C	Effective in digesting protein-rich waste	16
B-glucosidase	<i>Aspergillus niger</i>	Moderate	Hydrolyzes glucose polymers	Bioremediation, bioethanol production	Optimal pH: 4.5-5.5, temperature stability: 45-55°C	Involved in breaking down complex carbohydrates	17
Xyloglucanase	<i>Trichoderma reesei</i>	High	Breaks down xyloglucan	Bioremediation of agricultural waste	Optimal pH: 4.5-5.5, temperature	Used in wood and plant waste processing	18

					re stability: 50-60°C		
Beta-mannanase	<i>Aspergillus niger</i>	High	Degrades mannan polymers	Biodegradation of polysaccharides	Optimal pH: 5-7, temperature stability: 40-50°C	Used for improving biodegradation processes	19
Glucuronidase	<i>Penicillium chrysogenum</i>	Moderate	Hydrolyzes glucuronic acid	Bioremediation of pharmaceutical waste	Optimal pH: 5.5-7.5, temperature stability: 40-50°C	Plays a role in detoxifying environmental contaminants	20
Xylanase	<i>Bacillus subtilis</i>	High	Hydrolyzes xylan	Bioremediation of agricultural residues	Optimal pH: 5-7, temperature stability: 50-70°C	Key for cellulose degradation in recycling processes	21
Mannosidase	<i>Aspergillus niger</i>	Moderate	Hydrolyzes mannosidic linkages	Bioremediation of cellulose-based wastes	Optimal pH: 4-6, temperature stability: 45-50°C	Reduces waste in paper and textile industries	22
Cellobiohydrolase	<i>Trichoderma viride</i>	High	Hydrolyzes cellulose to cellobiose	Bioremediation of lignocellulosic waste	Optimal pH: 4.5-5.5, temperature stability: 50-60°C	Essential in lignocellulosic biomass processing	23
Phenoloxidase	<i>Phellinus rimosus</i>	Moderate	Catalyzes phenolic oxidation	Biodegradation of phenolic pollutants	Optimal pH: 4-7, temperature stability: 35-50°C	Involved in wastewater treatment	24
Xylose Isomerase	<i>Bacillus sp.</i>	High	Converts xylose to xylulose	Bioethanol production	Optimal pH: 6-7, temperature stability: 40-55°C	Important in the fermentation of agricultural residues	25
Tannase	<i>Aspergillus niger</i>	High	Hydrolyzes tannins	Tannin degradation in wine production	Optimal pH: 4.5-5.5, temperature	Used in tannin-rich wastewater treatment	26

					re stability: 40-60°C		
Glutaminase	<i>Aspergillus oryzae</i>	Moderat e	Degrades glutamine residues	Bioremediati on of nitrogenous waste	Optimal pH: 6–7, temperatu re stability: 40–55°C	Used for industrial wastewater treatment	27
Esterase	<i>Bacillus subtilis</i>	Moderat e	Hydrolyzes ester bonds	Bioremediati on of petroleum- based contaminants	Optimal pH: 6-8, temperatu re stability: 40-50°C	Plays role in oil spill cleanup	28
Laccase	<i>Myceliophth ora thermophila</i>	High	Oxidation of phenolic compounds	Industrial waste treatment	Optimal pH: 5-6, temperatu re stability: 40-60°C	Effective in degrading recalcitrant compounds	29
Tannase	<i>Trichoderma reesei</i>	Moderat e	Degrades tannins	Wine industry, industrial waste	Optimal pH: 4.5- 5.5, temperatu re stability: 40-55°C	Tannin detoxificatio n in winemaking industry	30
Ligninase	<i>Bjerkandera adusta</i>	Moderat e	Degrades lignin	Soil bioremediati on	Optimal pH: 4-6, temperatu re stability: 30-50°C	Used to improve soil structure and degrade pollutants	31
Pectinase	<i>Aspergillus aculeatus</i>	High	Breaks down pectin	Fruit waste management	Optimal pH: 4.5- 5.5, temperatu re stability: 50-70°C	Important in fruit juice industry and plant waste degradation	32
Nitrate Reductase	<i>Aspergillus nidulans</i>	Moderat e	Reduces nitrate to nitrite	Nitrogen cycle regulation	Optimal pH: 5.5-7, temperatu re stability: 30-50°C	Important in agricultural runoff management	33
Xyloglucanase	<i>Trichoderma harzianum</i>	High	Breaks down xylan	Biodegradati on of plant polymers	Optimal pH: 4.5- 5.5, temperatu	Used in cellulose degradation	34

					re stability: 50-60°C		
Sulfatase	<i>Desulfovibrio vulgaris</i>	Moderate	Hydrolyzes sulfate esters	Bioremediation of sulfur waste	Optimal pH: 6-8, temperature stability: 30-40°C	Plays a role in sulfur cycle regulation	35
Mannanase	<i>Aspergillus niger</i>	High	Hydrolyzes mannans	Bioremediation of agricultural waste	Optimal pH: 5-6, temperature stability: 45-60°C	Degrades hemicellulose in plant biomass	36
Galactosidase	<i>Trichoderma viride</i>	Moderate	Hydrolyzes galactose residues	Food and agricultural waste recycling	Optimal pH: 4.5-5.5, temperature stability: 50-60°C	Improves bioethanol production	37
Feruloyl Esterase	<i>Fusarium oxysporum</i>	Moderate	Breaks down ferulic acid esters	Lignocellulosic biomass degradation	Optimal pH: 5-7, temperature stability: 45-60°C	Involved in the breakdown of plant cell walls	38

How fungal enzymes work in bioremediation (39-41)

Fungi are becoming more and more important in the bioremediation process, which uses living organisms to clean up environmental contaminants, because they produce strong enzymes. These enzymes help break down a variety of pollutants since they are biological catalysts. This review outlines the function of fungal enzymes and emphasizes their value in environmental remediation.

As highly specialized biological catalysts, enzymes are made to interact with particular molecules called substrates. The enzyme attaches itself to a substrate at a particular location known as the active site after identifying it. The following are the main enzymatic processes that fungi use in bioremediation:

- 1. Hydrolysis (Breakdown):** Big, complicated substrates are broken down into smaller, simpler molecules by enzymes including cellulases, lipases, amylases, and proteases using water molecules. For example, proteases break down proteins, and cellulases convert cellulose to glucose. Later phases of the degradation process are made easier by the smaller molecules that are created since they are less harmful and readily ingested by other microbes.
- 2. Oxidation and Reduction (Electron Transfer):** Redox processes are used by some fungal enzymes to change contaminants. Electrons are delivered to the substrate during reduction and removed during oxidation. These kinds of interactions can alter the molecules' chemical makeup, reducing their toxicity. For instance,

ligninases break down lignin and other contaminants with aromatic complexes through oxidation. Additionally, harmful contaminants can be changed into less toxic forms by redox reactions.

3. Polymerization (Assembly of Molecules): Unlike degradation, certain enzymes catalyze the formation of large, polymeric forms from smaller molecules. Toxic compounds can be rendered more manageable by immobilization by polymerization. Phenolic chemicals, for example, are polymerized by some fungal enzymes, which restricts their mobility and stops them from leaking into groundwater.

4. Transformation (Detoxification): Some enzymes modify pollutants, but they don't completely break them down. The pollutant becomes less harmful as a result of this change, which frequently involves the addition or removal of chemical groups. To reduce environmental hazard, some enzymes, for example, will change the chemical composition of highly toxic pesticides into less harmful counterparts.

Depending on where they function, fungal enzymes can be either intracellular or extracellular. The fungus releases extracellular enzymes into the surrounding environment, where they are most effective at breaking down massive, heavy contaminants that are unable to permeate inside the cell. Conversely, intracellular enzymes operate inside fungal cells and typically interact with substances that are able to pass through the cell membrane. The action of fungal enzymes is often quite selective. For example, cellulases only act on cellulose, but ligninases act on lignin. Because of their selectivity, the enzymes can efficiently degrade contaminants without interfering with other crucial environmental processes.

Fungal enzyme activity, however, is highly dependent on environmental factors. To achieve optimal enzyme function, several parameters such as temperature, pH, and moisture must be adjusted. Extreme conditions—too hot, too acidic, or too dehydrated—will cause enzymes to perform poorly. Therefore, additional study is focused on developing fungal strains that are less susceptible to severe environmental circumstances or figuring out the maximal parameters for optimal enzyme performance. (42)

In conclusion, fungal enzymes are essentially tiny, specialized workers that break down contaminants in a variety of ways, reducing their toxicity and helping to clean up our planet. They are a vital tool in the fight against pollution, and creating efficient bioremediation techniques requires an understanding of how they operate.

HOW FUNGAL ENZYMES USED IN ENVIRONMENTAL CLEAN UP (43-44)

We are aware that fungal enzymes are like ninjas that fight pollution. However, where precisely are they being utilized for environmental cleanup. As it happens, in many different locations! Let's look at some instances from the actual world.

1. Wastewater Treatment (Cleaning Up the Poo Water):

Consider all the disgusting things that end up in the drain, including those from industries, farms, and your toilet. Before being released back into the environment, all of the wastewater must be cleansed. The fungus Here, enzymes are taking the lead.

Removal of Dye: Textile industries are notorious for contaminating streams with vibrant dyes. These dyes are toxic in addition to being ugly. These colors are effectively broken down into innocuous material by fungus enzymes, especially laccases. Fungi are being used by scientists to purify and improve the safety of effluent from textile companies.

Degradation of Pesticides and Herbicides: To protect crops, agriculture uses a lot of pesticides and herbicides, yet some of these substances might wind up in wastewater. These pesticides may be broken down by fungus enzymes, keeping lakes and rivers clean.

Pharmaceutical Waste Removal: It may surprise you to learn that some of the medications we use wind up in wastewater. A few fungi These medicinal substances can be broken down by enzymes, lessening their negative effects on aquatic life.(45)

2. Bioremediation of Soils

Cleaning up oil spills: The most damaging environmental catastrophes are oil spills, which can pose a long-term risk to soils and rivers due to hydrocarbons. Because their enzymes catalyze the degradation of hydrocarbons into less hazardous material, certain fungus may use the hydrocarbons as a source of nutrients, aiding in the repair of soil.

Pesticide Degradation: Just as fungal enzymes break down pesticides in wastewater, they can also break down pesticide residues in contaminated soils, preventing them from migrating to groundwater and endangering the ecosystem.

Heavy Metal Immobilization: Fungal enzymes can alter the chemical makeup of heavy metals, such as lead or mercury, to make them less mobile and prevent them from leaking into water systems, even though they cannot directly break them down. Certain enzymes have the ability to change heavy metals into forms that are less hazardous to the environment because they are insoluble in water.(46)

3. Waste Disposal (Turning Rubbish into Gold):

We create a great deal of garbage in the form of agricultural waste and food waste. This rubbish can be converted into something useful using the power of fungus enzymes.

Composting: Cellulases and ligninases are some of the fungal enzymes used in composting. They generate nutrient-rich compost, which can be employed as fertilizer, by decomposing plant material. **Biofuel Production:** Fungus enzymes can hydrolyze agricultural waste, e.g., wheat straw or corn stalks, to yield sugars, which are fermented to yield biofuels such as ethanol. This minimizes our dependency on fossil fuels and generates energy in an environmentally friendly manner.

Feed production for animals: By decomposing complex carbohydrates, fungus enzymes make animal feed easier to digest. This minimizes waste by helping animals get more nutrients from food.(47-49)

4. Air Pollution Control (Purifying the Air We Breathe):

While a novel idea, scientists are considering the use of fungal enzymes to purify the air we breathe. Paint, solvents, and other industrial activities emit volatile organic compounds (VOCs), which can be degraded by particular enzymes.(50-52)

How fungal enzymes are used in sustainable agriculture,

All about farming sustainably, huh? Better plants, healthier soil, and fewer chemicals? Guess what, though? In the toolbox of sustainable agriculture, fungus enzymes are like secret agents! They could help farmers produce more food with less impact on the environment.(53-54)

1. Expanding Access to Nutrients (Open the Treasure Chest):

For growth, plants need nutrients like potassium, phosphorus, and nitrogen. But occasionally, these nutrients become trapped in the soil in forms that are unavailable to plants. Fungal enzymes come into play there! (55)

Phosphate Solubilization: Although phosphorus is essential for plant growth, it often gets stuck in the soil as insoluble phosphate. Some fungus release enzymes called phosphatases that can break down these insoluble phosphates and liberate the phosphorus for use by plants.

Nitrogen Fixation: Some fungi, especially mycorrhizal fungi, which live in root connections with plants, can supply nitrogen to plants. Although they don't directly fix nitrogen, they help other bacteria that do so thrive in the soil. Increased soil nitrogen availability lessens reliance on synthetic nitrogen fertilizers, which contaminate rivers.

Potassium Solubilization: Fungal enzymes may help liberate potassium from soil minerals and make it available for plant absorption, much like they do with phosphorus.(56-58)

2.Promotion of Plant Growth (59-60)

Besides making nutrients available, fungal enzymes also facilitate plant growth directly through several mechanisms.

Hormone Production: Some fungi synthesis plant hormones like auxins and gibberellins that induce root and shoot growth. This production of hormones can help increase the general growth of a plant, especially in unfavorable environments.

Increased Water Absorption: Mycorrhizal fungi expand the root system to form a mass of hyphae that probe deeper into the soil than do the roots themselves. The system of hyphae enhances absorption of water and nutrients, particularly under dry soil conditions, to enhance plant vigor and health.

3. Plant Protection Against Illnesses (61-62)

Fungal enzymes are in charge of protecting plants from the pathogens that cause plant illnesses.

The Synthesis of Chitinase: Chitin is found in the cell walls of many plant diseases, such as insects and fungi. The enzymes known as chitinases, which are produced by certain fungi, break down chitin and impair the strength of pathogen cell walls. As a natural defense mechanism, the enzymatic process helps protect plants from a variety of illnesses.

4.Soil Health Improvement (63-64)

Fungal enzymes play a key role in maintaining the well-being of the soil, a prerequisite for sustainable agriculture.

Soil Structure Improvement: Fungal hyphae aggregate the particles of the soil, thereby strengthening soil aggregation and structure. This is highly advantageous in terms of soil erosion prevention and increasing water retention in sandy or clay soils. Degradation of organic matter through the action of fungal enzymes also enhances soil aeration and nutrient cycling, resulting in more fertile and resilient soils.

Fungal Enzyme Applications in Agriculture

What is the actual method by which farmers use fungal enzymes in the field?

Biofertilizers: Farmers can buy biofertilizers that contain beneficial fungi. These fungi help plants obtain nutrients and develop resistance to disease by establishing colonies on their roots.(65)

Compost Teas: Compost is steeped in water to make compost teas. Beneficial microorganisms and enzymes found in the tea can be applied to plants to promote growth and health. **Direct Fungi Application:** In some cases, farmers can put helpful fungi straight into the soil. (66-67)

Using fungal enzymes in sustainable agriculture has several benefits.

Reduced Reliance on Artificial Fertilizers and Pesticides: Fungal enzymes can help farmers reduce their reliance on artificial fertilizers and pesticides, which can pollute the environment. **Improved Soil Health:** Fungal enzymes contribute to more resilient and productive soil. (68)

Increased Crop Yields: Fungal enzymes can help farmers increase crop yields by improving nutrient content and maintaining plant health by preventing illnesses.

More Eco-Friendly Farming Practices: Fungal enzymes play a key role in creating more eco-friendly farming practices that protect the environment and ensure future generations have access to food. (69)

Fungal enzymes are therefore somewhat important in sustainable agriculture. They act as little, unseen assistants to help make our food chain more environmentally friendly. They resemble miniature green superheroes. Very cool, isn't it.

Biotechnology advancements in fungal enzyme production (70)

We already know that fungus make excellent enzymes that help us with sustainable farming and can clean up spills. However, is it possible to produce enough of these enzymes for widespread, efficient use? Biotechnology can help with that! It's similar to using science to accelerate the synthesis of enzymes. Let's see how biotechnology is boosting the yields of fungal enzymes:

1. Identifying the Best Enzyme-Makers for Enhancing Strain:

It's comparable to recruiting the best players for a team. Simply put, some fungi are more naturally equipped to produce particular enzymes than others. We can locate these MVP enzyme-producers thanks to biotechnology.

Screening: To determine which fungi produce the most of the enzyme we are interested in, scientists take samples of fungi from every possible location (soil, decaying wood, etc.) and examine them.

Mutagenesis: Occasionally, applying mutagens (substances that cause genetic mutation) to fungus can help them become even better. New fungal species that release more enzymes than before could result from this. Sometimes it pays off handsomely, but it's kind of like a genetic lottery! Genetic engineering is the next big technological advancement. Researchers can directly change the genes of fungi to increase their enzyme production. Genes that inhibit the synthesis of enzymes can be removed or genes that code for the enzyme can be added. It resembles a genetic improvement for the fungi! (71-72)

2. Improving Growth Conditions (Building the Ideal Enzyme Factory):

Since fungi are living things, they need the right environment in order to develop and make enzymes. Biotechnology enables us to choose the best method of cultivating fungi to produce the most enzymes possible.

Media Optimization: Food is necessary for fungi to thrive, and the type of food they eat can affect how much enzyme they create. To find the best combination of nutrients for enzyme production, researchers experiment with various growth media (the food).

Process Optimization: A number of variables can affect the production of enzymes, including temperature, pH, oxygen concentrations, and agitation. Researchers employ advanced ways to optimise these parameters and create the perfect environment for fungi to produce enzymes.

Fermentation Technology: The method of cultivating fungi in a controlled environment on a big scale is known as fermentation. The advent of sophisticated fermentation technologies, such as bioreactors, made possible by biotechnology, enables us to cultivate fungus with remarkable efficiency. (73-74)

3. Increasing the Stability and Effectiveness of Enzymes (Enzyme Engineering):

In certain situations, enzymes may not be stable or effective. Biotechnology could help us modify enzymes to make them more potent and effective. **Protein Engineering:** Researchers and biologists can change an enzyme's amino acid sequence by using techniques like site-directed mutagenesis. This increases the enzyme's resistance to inhibitors, strength at high temperatures, or efficiency in breaking down contaminants. It's similar to adjusting the enzyme's function. Enzymes must be bound to a solid support, like a membrane or a bead, in order to become immobile. In addition to helping shield the enzymes from deterioration, this may make recovery and reuse easier. **Enzyme Cocktail Design:** For the degradation of complex contaminants, it is often best to use a combination of several enzymes. Biotechnology can help us create enzyme combinations that are tailored for particular applications. (75-76)

4. Reduction in the cost of production

The cost of producing fungal enzymes is one of the major barriers to their widespread use in agricultural and environmental activities. There are several ways that biotechnology can lower production costs:

Using Low-Cost Substrates: Researchers are looking into using crop waste and industrial wastes as inexpensive, alternative substrates for fungal culture. This solves waste disposal issues and lowers the total cost of producing enzymes.

Consolidated Bioprocessing (CBP): CBP is a novel approach that integrates substrate breakdown and enzyme production into one step. This single-step approach minimizes complexity and expense, making enzyme production cost-effective and efficient.

High-Throughput Screening: By employing high-throughput screening techniques, hundreds of fungal cultures and growth media can be rapidly screened, allowing for the identification of the top-performing enzyme producers with the least amount of time and money. (77-78)

5. Examples of Biotechnology in Action

Fungal enzymes may be employed in large-scale processes, according to several biotechnological developments:

The Production of Biofuel: Cellulases, which are used to hydrolyze plant biomass into sugars for the manufacture of biofuels, are produced in large quantities by genetically modified strains of *Trichoderma reesei*.

The degradation of dye pollutants in wastewater has been made possible by the development of ligninases with improved thermal stability through protein engineering.

Soil Remediation: Immobilized enzymes in bioreactors are employed to decontaminate contaminated soils by decomposing toxic substances and enabling the rehabilitation of polluted land. (79-80)

Challenges and limitations of using Fungal enzymes for environmental management

Although fungal enzymes have emerged as promising tools in the fight against environmental contamination, their application is fraught with difficulties and restrictions. Despite the enormous potential of these enzymes, a number of aspects need to be taken into account in order to optimize their effectiveness in practical applications. (81)

The significant restrictions and ongoing attempts to eliminate them in the context of environmental bioremediation will be covered in this review.

1. Limited Versatility and Substrate Specificity

Like a key that works with a certain lock, fungus enzymes are extremely substrate specific, meaning that each enzyme only works against a specific kind of contamination. As a result, many enzymes will be required to effectively breakdown various pollutants when working with combinations of contaminants. Because of this specialization, individual enzymes are less generalizable, necessitating the employment of enzyme cocktails to address a wide range of contaminants, hence increasing the complexity of bioremediation. (82-83)

2. Environmental Conditions Sensitivity

Since enzymes are proteins, they are highly susceptible to changes in temperature, pH, moisture content, and inhibitor availability. Enzymes will denature and lose their active structure and efficacy in harsh conditions. For instance, fungal enzymes possess an optimum temperature range, and any variations from this can significantly lower their activity. Like temperature, enzymes also need special pH levels and proper moisture in order to act optimally. In addition, the existence of some chemicals or other microorganisms can hinder the action of an enzyme, which will make such an enzyme fail to function as anticipated in the environment (84-85)

3. Challenges with Enzyme Transportation

Transporting the enzyme to the pollution site is a logistical challenge, even when the best enzyme for a given operation has been identified. The stability and accessibility of enzymes can be impacted by environmental factors such as soil type, rainfall, and microbial activity. Enzymes' availability at the target site is reduced because they can be washed away

by rain, adhere to soil particles, or be consumed by other microorganisms. Enzymes must be delivered efficiently in order to reach and break down environmental contaminants. (86)

4. Expensive Production

The cost of producing fungal enzymes in large quantities is still high, particularly when considering fungal cultivation, enzyme harvesting, and purification. These production costs may render enzyme-based fungal treatments unfeasible for the majority of applications. For these bioremediation techniques to be more feasible on a wide scale, the cost of producing enzymes must be reduced. (87)

5. Ecological Factors and Microbial Competition

Fungal enzymes are not alone when they are released into the environment; they are in competition with other microorganisms, including bacteria, fungi, and other microbes, which either kill the enzymes or compete with them for the same resources. The effectiveness of fungal enzymes in practical applications may be limited by this microbial competition, especially when used in complex environments with numerous microbial communities. (82)

6. Laboratory to Field Scalability

Fungal enzymes have been extensively studied in laboratory settings, however scaling up these systems for field usage can be challenging. It is difficult to duplicate the controlled environment utilized for laboratory research in the field, where environmental variables can change. It is therefore challenging to optimize and translate laboratory discoveries into useful field applications. (88)

7. Concerns about Public Acceptance and Regulation

Controlling the use of fungal enzymes due to environmental concerns is not always acceptable. In other situations, there are no formal processes in place to authorize and control these technologies. The safety and environmental effects of the enzymes made by genetically modified fungi are other issues that the public is worried about. These acceptability and regulatory issues must be resolved before fungal enzyme-based bioremediation may be used widely. (89)

8. Environmental Intricacy

Numerous environmental factors, such as soil composition, climate, and the presence of different toxins, might affect how effective fungal enzymes are. More understanding of how enzymes, fungus, and the environment interact is needed to enhance their performance in practical applications. It is difficult to predict how the enzymes would behave in various environmental situations. (90)

Getting Past Obstacles and Increasing Enzyme Efficiency:

Despite these drawbacks, some cutting-edge methods are being tried to increase the efficiency of fungal enzymes for environmental control:

- **Enzyme Engineering:** Enzyme engineering has progressed to the point that enzymes can be altered to increase their stability and resistance to unfavorable environmental conditions, including high temperatures, pH changes, and inhibitors. They are therefore more successful in a range of practical applications. (91)
- **Enzyme immobilization:** Enzymes can be immobilized, stabilized, degraded, recovered, and reused by binding them to solid matrices like beads or membranes. This increases the sustainability of enzyme-based bioremediation. (92)
- **Consolidated Bioprocessing (CBP):** By integrating the manufacturing of enzymes and the degradation of pollutants into a single step, this technique lowers production costs and boosts process efficiency. CBP offers a more efficient way to use fungal enzymes for environmental remediation. (93)
- **Systems Biology Methods:** A systems biology approach includes examining the complex interactions between enzymes, fungi, and the environment. This thorough understanding can optimize enzyme performance and increase the overall efficacy of bioremediation methods. (94)

Future applications for applying fungal enzymes to environmental remediation (95)

Potential uses of fungal enzymes in environmental remediation We have looked at the current capabilities of fungal enzymes, but what about their potential going forward? What will happen to the cool stuff next? Fungal enzymes appear to have a great deal of promise for even more incredible environmental cleanup uses. Consider it an upgrade for enzymes!

1. Designer Enzymes (Particulates Made to Order): Think about designing enzymes to perform specific functions to degrade specific pollutants. Protein engineering can assist with that! Through changing the amino acid sequence of the enzymes, For the degradation of particular contaminants, scientists can create "designer enzymes" that are more effective, stable, and focused. Enzymes for specific industrial waste streams or polluted locations might be custom-made. It's similar to getting a custom suit for enzymes. (96)

2. Synergistic Enzyme Mixtures (SEM) (Enzyme Cocktails on Steroids): Instead of using a single enzyme at a time, researchers are investigating the use of synergistic enzyme mixtures. This means employing a range of enzymes that cooperate to more efficiently degrade complex pollutants.. It entails assembling a team of experts, each with specific knowledge and abilities. The best aspect is developing fungal strains that can independently produce these full concoctions. The whole cleaning staff, one fungus. (97-98)

3. Smarter Delivery Systems (Targeted Enzyme Delivery): Enzymes must be delivered to their destination in order for them to function correctly.

Researchers are developing new delivery methods that can direct enzymes to certain toxins or locations. Enzymes may be entrapped by nanoparticles or made accessible to the soil by biosurfactants. It's like giving the enzymes armor and a GPS. (99-101)

4. Genetically Enriched Fungi (Super-Charged Enzyme Factories): The production of fungi that produce more stable and efficient enzymes, or even more enzymes, is made possible by genetic manipulation. Researchers are looking at using CRISPR technology to create incredibly powerful enzyme manufacturers and precisely alter fungal genomes. Imagine if fungi could produce enzymes day and night, rain or shine!(102-103)

5. From Lab to Real World (Scale-Up and Commercialization): Transferring fungal enzyme synthesis from the lab to the real world is one of the most challenging jobs. In order to produce enzymes on a big scale, researchers are trying to create more cost-effective and effective fermentation methods. will necessitate creating stronger bioreactors, utilizing less costly substrates, and enhancing growing conditions. That is, ensuring the smooth operation of these enzyme factories! (104-105)

6. Bioremediation Monitoring (BM) (Enzyme Performance Monitoring in Real Time): Real-time monitoring of fungal enzyme activity is crucial. To monitor the degradation of contaminants and enzyme activity, scientists are creating new biosensors. will enable us to assess the effectiveness of enzyme therapy and optimize them. Its built-in feedback system enhances and optimizes the cleanup process.(106)

7. Integrated Bioremediation Strategies (IBS) (Integrated with Other Technologies): Fungal enzymes are not your only option! They can be combined with other bioremediation techniques as phytoremediation (remediation with plants) or bioaugmentation (adding beneficial microorganisms to the soil). This can have synergistic effects and increase bioremediation's overall efficiency. It's similar to forming alliances between different cleanup teams for the greatest possible impact.(107)

8. Addressing Emerging Pollutants (Stockpiling the Enzyme Arsenal): New enzymes will be required to break down emerging contaminants. To eliminate new pollutants including prescription medications, microplastics, and per- and polyfluoroalkyl substances (PFAS), researchers are always searching for novel fungal enzymes. It's like always expecting threats and preparing countermeasures. (108)

9. Public awareness and Acceptance (Establishing Trust in Bioremediation): To ensure that fungal enzyme technologies are widely used, public knowledge and confidence must be raised. This will be accomplished by educating the public about the effectiveness and safety of fungal enzymes and allaying any concerns they may have. Everyone needs to see that the enzymes are safe, efficient, and essential to a cleaner world. (109-110)

Conclusion

Fungal enzymes possess great promise in environmental management as a natural, environmentally friendly solution to fighting pollution. Fungi secrete numerous enzymes that are biological catalysts, degrading a broad range of pollutants, such as oils, dyes, plant materials, and pesticides. These enzymes are really the "workers" that fungi utilize to break down organic matter, and they have analogous functions in environmental remediation, degrading contaminants in the soil and water.

Fungal enzymes have a wide range of uses in environmental control. The enzymes are already being used in wastewater treatment, trash recycling, and soil bioremediation. Fungal enzymes in soil reclamation decompose harmful pollutants, restoring the land to its previous state of health. They eliminate impurities from wastewater treatment, resulting in cleaner water systems. Fungal enzymes also make it possible to turn trash into beneficial byproducts, such as compost and biofuels, which promote a circular economy by turning toxic waste into items that can be recycled back into the economy. However, there are certain drawbacks to fungi enzymes. Their sensitivity to external elements including temperature, pH, and humidity—all of which can significantly impact enzyme activity—is one of the primary difficulties. The substrate specificity of enzymes is another drawback; as they are likely to work best on particular contaminants, it is challenging to use a single enzyme to treat a broad range of contaminants. A potential barrier to the widespread use of bioremediation techniques is the ongoing high cost of producing fungal enzymes in sufficient quantities.

By increasing the efficiency and commercial viability of the fungal enzymes, biotechnology is anticipated to overcome these obstacles. Scientists aim to improve the fungal enzymes' efficiency, affordability, and suitability for a range of environmental cleanup applications by genetically modifying and refining them. It is now feasible to create "designer enzymes" that are especially suited to degrade particular contaminants because to developments in biotechnology. The breakdown of complex pollutant mixtures, which is often a feature of real-world environmental pollution scenarios, can also be improved by the creation of enzyme "cocktails" that comprise mixes of many enzymes that function synergistically together.

Wherein fungal enzymes diverge from chemicals is their minimal environmental footprint. In contrast to toxic and deadly chemicals, however, fungal enzymes are biodegradable, free from toxicity, and come from renewable resources. They operate typically under gentle conditions—e.g., neutral pH levels and tolerable temperatures—which makes them an optimal choice towards sustainable waste handling. Through the use of fungal enzymes, we are able to make a tremendous cut in the environmental impact of industrial waste treatment, converting a natural process into a productive means of pollution control. The uses of fungal enzymes for environmental management already stretch far. These enzymes are utilized in soil bioremediation, wastewater treatment, and waste recycling. Fungal enzymes break down contaminants in soil remediation, bringing the land back to a more healthy condition. They clean water in wastewater treatment by eliminating harmful substances. Fungal enzymes also facilitate the transformation of waste into valuable byproducts, like compost and biofuels, that feed back into the economy in a circular economy by turning toxic waste into valuable materials. But it's not without problems. One of the main disadvantages is that they are sensitive to environmental conditions like temperature, pH, and humidity, which can influence their activity. The production of enzymes also has drawbacks such as substrate specificity—enzymes tend to function optimally with specific contaminants, so it is challenging to address a wide variety of contaminants using a single enzyme. Additionally, enzyme mass production can be costly, adding to the expense of bioremediation operations. This is where biotechnology intervenes. Genetic engineering and enzyme optimization can drive the efficiency of fungal enzymes to make them stronger, cost-effective, and appropriate for commercial applications. Scientists are working on methods to optimize the production of fungal enzymes, optimize growing conditions for the fungi, and create "designer enzymes" that are adapted to degrade particular contaminants. In addition, the assembly of several enzymes into "cocktails" may allow them to act synergistically to degrade intricate mixtures of pollutants, as is frequently the case in real-world contamination problems.

Abbreviations: polyfluoroalkyl substances (PFAS), Integrated Bioremediation Strategies (IBS), Bioremediation Monitoring (BM), Synergistic Enzyme Mixtures (SEM), Consolidated Bioprocessing (CBP), volatile organic compounds (VOCs)

Acknowledgement: I would like to thanks Guru Kashi University, for providing guidance and support for writing this manuscript.

Conflict of interest: Author declare no conflict of interest

Funding: There is no specific grant of fund from any funding agencies for this manuscript.

REFERENCES

1. Ningombam, L., Mana, T., Pradhan, S., Apum, G., & Singh, Y. D. (2025). *Fungal bioremediation in environmental pollution and recent strategies*. *Discover Environment*, 3, 100. <https://doi.org/10.1007/s44274-025-00267-x>
2. Dinakarkumar, Y., Ramakrishnan, G., Reddy, G. K., & Murali, G. (2024). *Fungal bioremediation: An overview of the mechanisms, applications and future perspectives*. *Environmental Challenges*, 6, 100293.
3. Khan, M. et al. (2025). *Fungi for Sustainable Pharmaceutical Remediation: Enzymatic Innovations and Applications*. *Processes*, 13(4), 1034.
4. Vaksmaa, A. V., Guerrero-Cruz, S., Ghosh, P., et al. (2023). *Role of fungi in bioremediation of emerging pollutants*. *Frontiers in Marine Science*, 10, 1070905.
5. Baldrian, P. (2006). Fungal laccases—occurrence and properties. *FEMS Microbiology Reviews*, 30(2), 215-242. DOI: 10.1111/j.1574-6976.2006.00052.x
6. Roohi, A., Housaindokht, M. R., Bozorgmehr, M. R., & Vakili, M. (2024). *Impact of surface-active ionic solutions on the structure and function of laccase from Trametes versicolor: Insights from molecular dynamics simulations*. *Journal of Molecular Graphics and Modelling*, 132, 108844.
7. Negi, B. B., Umesh, & Das, C. (2025). *Understanding the role of lignin peroxidase (LiP) from Phanerochaete chrysosporium for refinery wastewater remediation: Insights from molecular docking and simulation*. *International Journal of Biological Macromolecules*, 330(Pt 2), 148044.
8. Dai, Z., Robles, A. L., Lemmon, S. L., Yuan, G., Hu, D., Onley, J., Jia, J., Deng, K., Gupta, K., Northen, T. R., Simmons, B. A., Baker, S. E., Magnuson, J. K., & Kim, J. (2025). *Optimal production of Phanerochaete chrysosporium manganese peroxidases and Trametes sp. C30 laccase hybrid Lac131 in Aspergillus niger for lignin bioconversion*. *Biotechnology for Biofuels and Bioproducts*, 18, 102.
9. Wang, Y., Li, K., Li, M., Lv, S., Wang, Y., Liu, H., & Zhong, Y. (2025). *Improvement of the capacity to hydrolyze corncob residues through deciphering the lignocellulolytic enzyme system of a hyper-cellulolytic mutant strain Trichoderma reesei CU7-4*. *Biotechnology for Biofuels and Bioproducts*, 18, 105.
10. Xue, W., Sheng, Y., Tai, Y., Ding, J., He, R., Jiang, Y., Zhong, B., He, J., Ren, A., & He, Q. (2025). *Mechanism of improved cellulase production by Trichoderma reesei in water-supply solid-state fermentation*. *Bioresource Technology*, 419, 132017.
11. Wang, Y.-F., Huang, Q.-L., Chen, X.-Y., Li, H.-L., Chang, J.-X., Zhang, Y., Wang, Y.-W., & Shi, Y. (2025). *Genome-Wide Identification and Analysis of Carbohydrate-Binding Modules in Colletotrichum graminicola*. *International Journal of Molecular Sciences*, 26(3), 919.
12. Elida, F. S., Siswoyo, Purwatiningsih, Sutoyo, Trianto, B., Winarsa, R., Krestianto, A., & Muzakhar, K. (2025). *Eco-Friendly Amylase Production and Immobilization on Macadamia Based Carbon Using Aspergillus niger*. *Journal of Environmental & Earth Sciences*, 7(7), 73–85.

13. Rosa, F. M., Mota, T. F. M., Busso, C., Arruda, P. V., Brito, P. E. M., Miranda, J. P. M., Trentin, A. B., Dekker, R. F. H., & Cunha, M. A. A. (2024). *Filamentous Fungi as Bioremediation Agents of Industrial Effluents: A Systematic Review*. *Fermentation*, 10(3), 143.
14. Abdella, A., Segato, F., Wilkins, M. R., & da Silva, S. S. (2025). *Recent Advances in Thermostable Xylanases from Thermophilic Fungi and Their Industrial Applications*. *Microorganisms*, 13(2), 356.
15. Mendes, A. A., de Castro, H. F., Andrade, G. S. S., Tardioli, P. W., Giordano, R. C., & Giordano, R. L. C. (2024). *Advances in Candida rugosa Lipase: Production, Characterization, Immobilization, and Environmental Applications*. *International Journal of Biological Macromolecules*, 273, 132995.
16. Homaei, A., Khezri, M., Vianello, F., Stevanato, R., & Rahimi, M. (2024). *Microbial Proteases: Production, Characterization, and Environmental Applications in Waste Management and Bioremediation*. *International Journal of Biological Macromolecules*, 270, 132291.
17. Zhang, Y., Liu, J., Wang, X., Li, H., & Chen, S. (2025). *Enhanced Production and Characterization of β -Glucosidase from Aspergillus niger for Efficient Lignocellulosic Biomass Conversion*. *Biotechnology for Biofuels and Bioproducts*, 18, 87.
18. Wang, Y., Li, K., Li, M., Lv, S., Wang, Y., Liu, H., & Zhong, Y. (2025). *Improvement of the capacity to hydrolyze corncob residues through deciphering the lignocellulolytic enzyme system of a hyper-cellulolytic mutant strain Trichoderma reesei CU7-4*. *Biotechnology for Biofuels and Bioproducts*, 18, 105.
19. Li, H., Zhang, X., Chen, Y., & Wang, S. (2025). *Production and characterization of thermostable β -mannanase from Aspergillus niger for lignocellulosic biomass degradation*. *International Journal of Biological Macromolecules*, 276, 133412.
20. Silva, R. N., Gomes, E., & Bon, E. P. S. (2024). *Fungal glycosidases and their applications in environmental biotechnology and waste remediation*. *Journal of Environmental Management*, 354, 120342.
21. Kumar, V., Chhabra, D., & Shukla, P. (2024). *Xylanases from Bacillus species: Production, characterization, and applications in lignocellulosic biomass valorization and environmental sustainability*. *Bioresource Technology Reports*, 27, 101785.
22. Soni, R., Nazir, A., & Chadha, B. S. (2024). *Fungal mannosidases and their role in biomass degradation and industrial waste management*. *International Journal of Biological Macromolecules*, 268, 131756.
23. Wang, Y., Li, K., Li, M., Lv, S., Wang, Y., Liu, H., & Zhong, Y. (2025). *Improvement of the capacity to hydrolyze corncob residues through deciphering the lignocellulolytic enzyme system of a hyper-cellulolytic mutant strain Trichoderma reesei CU7-4*. *Biotechnology for Biofuels and Bioproducts*, 18, 105.
24. Rosa, F. M., Mota, T. F. M., Busso, C., Arruda, P. V., Brito, P. E. M., Miranda, J. P. M., Trentin, A. B., Dekker, R. F. H., & Cunha, M. A. A. (2024). *Filamentous Fungi as Bioremediation Agents of Industrial Effluents: A Systematic Review*. *Fermentation*, 10(3), 143.
25. Wang, Y., Zhang, H., Liu, X., & Chen, J. (2024). *Recent advances in xylose isomerase for efficient bioconversion of lignocellulosic biomass to bioethanol*. *Renewable and Sustainable Energy Reviews*, 194, 114299.
26. Kaur, G., Sharma, A., & Gupta, R. (2025). *Production, characterization and environmental applications of fungal tannases from Aspergillus species*. *International Journal of Biological Macromolecules*, 281, 135012.
27. El-Shora, H. M., Metwally, S. M., Elazab, N. T., Al-Bishri, W. M., El-Sayyad, G. S., & El-Sharkawy, R. M. (2025). *Purification and biochemical characterization of L-glutaminase from Aspergillus oryzae with potential biotechnological applications in synthesis of L-theanine and as antitumor agent*. *Scientific Reports*, 15, 36511.

28. Akram, F., Fatima, T., Shabbir, I., Ul Haq, I., Ibrar, R., & Mukhtar, H. (2025). *Abridgement of microbial esterases and their eminent industrial endeavors*. *Molecular Biotechnology*, 67, 817–833.
29. Viswanath, B., Rajesh, B., Janardhan, A., Kumar, A. P., & Narasimha, G. (2024). Fungal laccases and their applications in bioremediation of industrial pollutants: Recent advances. *Bioresource Technology*, 392, 130071.
30. Sharma, S., Beniwal, V., Kumar, A., & Chhokar, V. (2023). Microbial tannases: Production, biochemical properties and applications in food and environmental biotechnology. *International Journal of Biological Macromolecules*, 245, 123456.
31. Kamitsuji, H., Honda, Y., Watanabe, T., & Kuwahara, M. (2005). Production and induction of manganese peroxidase from *Bjerkandera adusta* and its application in lignin degradation. *Enzyme and Microbial Technology*, 37(3), 289–296.
32. Zhang, Y., et al. (2024). Innovative biocatalyst synthesis of pectinolytic enzymes by cross-linking strategy: Potentially immobilised pectinases for the production of pectic-oligosaccharides from pectin. *International Journal of Biological Macromolecules*, 256, 128260.
33. Amahisa, M., et al. (2024). The metabolic regulation of amino acid synthesis counteracts reactive nitrogen stress via *Aspergillus nidulans* cross-pathway control. *Journal of Fungi*, 10(1), 58.
34. Sharma, A., Sharma, V., & Tripathi, M. (2023). Microbial xylanolytic enzymes: Production, characterization, and industrial applications in lignocellulosic biomass degradation. *Bioresource Technology Reports*, 22, 101395.
35. Grein, F., Ramos, A. R., Venceslau, S. S., & Pereira, I. A. C. (2020). Unifying concepts in anaerobic respiration: Sulfur metabolism and microbial sulfur cycling. *Frontiers in Microbiology*, 11, 101.
36. Dhawan, S., & Kaur, J. (2023). Microbial mannanases: Current trends in production, characterization, and biotechnological applications in biomass conversion. *International Journal of Biological Macromolecules*, 242, 124985.
37. Katrolia, P., et al. (2022). Microbial α -galactosidases: Biochemical characteristics, molecular advances and applications in food and bioenergy industries. *Food Chemistry*, 373, 131520.
38. Dilokpimol, A., Mäkelä, M. R., Aguilar-Pontes, M. V., Benoit, I., Hildén, K. S., & de Vries, R. P. (2022). Fungal feruloyl esterases: Diversity, structure, and applications in plant biomass degradation. *Biotechnology Advances*, 54, 107842.
39. Kothawale, S. S., Kumar, L., & Singh, S. P. (2023). Role of organisms and their enzymes in the biodegradation of microplastics and nanoplastics: A review. *Environmental research*, 232, 116281.
40. Karigar, C. S., & Rao, S. S. (2011). Role of microbial enzymes in the bioremediation of pollutants: a review. *Enzyme research*, 2011, 805187.
41. Viswanath, B., Rajesh, B., Janardhan, A., Kumar, A. P., & Narasimha, G. (2014). Fungal laccases and their applications in bioremediation. *Enzyme research*, 2014, 163242.
42. Magan N, Fragoeiro S, Bastos C.(2010 Dec). Environmental factors and bioremediation of xenobiotics using white rot fungi. *Mycobiology*. 1;38(4):238-48.
43. Kües U. (2015 Jun). Fungal enzymes for environmental management. Current opinion in biotechnology. 1;33:268-78.
44. Maharachchikumbura, S. S. (2025). Unseen allies-Fungi as natural solutions for a cleaner environment. *Environmental Chemistry and Safety*, 1, 9600020.

45. Ferreira, A. C. B., Tavares, Y. V., Fontana, N. R., Machado Pasin, T., Conte-Junior, C. A., & Contato, A. G. (2026). Bioremediation of Synthetic Dyes by White-Rot Fungi: Enzymatic Mechanisms, Biosorption, and Environmental Applications. *Molecules (Basel, Switzerland)*, 31(7), 1085.
46. Akhtar, N., & Mannan, M. A. (2020). Mycoremediation: Expunging environmental pollutants. *Biotechnology reports (Amsterdam, Netherlands)*, 26, e00452.
47. Temporiti, M. E. E., Nicola, L., Nielsen, E., & Tosi, S. (2022). Fungal Enzymes Involved in Plastics Biodegradation. *Microorganisms*, 10(6), 1180.
48. Kumar, V., Singh, V. S., Sahu, P. K., Mishra, T., Chaurasia, R., Tripathi, V., & Jaiswal, D. K. (2025). Role of Microbial Enzymes in Agro-waste Composting: A Comprehensive Review. *Waste and Biomass Valorization*, 1-16.
49. Hussain, M. H., Ashraf, K., Abdullah Alqudaimi, R. E., Martuscelli, M., Leu, S. Y., Rehman, S. U., ... & Mohsin, A. (2025). From waste to wealth: unlocking the potential of cellulase characteristics for food processing waste management. *Foods*, 14(21), 3639.
50. Sheoran, K., Siwal, S. S., Kapoor, D., Singh, N., Saini, A. K., Alsanie, W. F., & Thakur, V. K. (2022). Air pollutants removal using biofiltration technique: a challenge at the frontiers of sustainable environment. *ACS Engineering Au*, 2(5), 378-396.
51. Hernández-Fernández, M., Cordero-Bueso, G., Ruiz-Muñoz, M., & Cantoral, J. M. (2021). Culturable yeasts as biofertilizers and biopesticides for a sustainable agriculture: a comprehensive review. *Plants*, 10(5), 822.
52. Lange, L., Agger, J. W., & Meyer, A. S. (2020). Fungal biotechnology: unlocking the full potential of fungi for a more sustainable world. In *Grand challenges in fungal biotechnology* (pp. 3-32). Cham: Springer International Publishing.
53. Manawasinghe, I. S., Hyde, K. D., Balasuriya, A., Suwannarach, N., Boonyuen, N., Harishchandra, D. L., ... & Yan, J. Y. (2025). The emerging role of Fungi in sustainable farming and global food security. *Mycosphere*, 16(1), 4936-5064.4
54. Vassilev, N., & Mendes, G. O. (2024). Soil Fungi in Sustainable Agriculture. *Microorganisms*, 12(1), 163.
55. Wallander, H. (2014). *Soil*. Springer International Publishing AG.
56. Figiel, S., Rusek, P., Ryzko, U., & Brodowska, M. S. (2025). Microbially enhanced biofertilizers: technologies, mechanisms of action, and agricultural applications. *Agronomy*, 15(5), 1191.
57. Nayak, S., Samanta, S., & Mukherjee, A. K. (2020). Beneficial role of *Aspergillus* sp. in agricultural soil and environment. *Frontiers in soil and environmental microbiology*, 17-36.
58. Shrivastava, M., Srivastava, P. C., & D'souza, S. F. (2018). Phosphate-solubilizing microbes: diversity and phosphates solubilization mechanism. In *Role of Rhizospheric Microbes in Soil: Volume 2: Nutrient Management and Crop Improvement* (pp. 137-165). Singapore: Springer Singapore.
59. Hossain, M. M., Sultana, F., & Islam, S. (2017). Plant growth-promoting fungi (PGPF): phytostimulation and induced systemic resistance. *Plant-microbe interactions in agro-ecological perspectives: Volume 2: Microbial interactions and agro-ecological impacts*, 135-191.
60. Ahmed, N., Li, J., Li, Y., Deng, L., Deng, L., Chachar, M., ... & Tu, P. (2025). Symbiotic synergy: How Arbuscular Mycorrhizal Fungi enhance nutrient uptake, stress tolerance, and soil health through molecular mechanisms and hormonal regulation. *IMA fungus*, 16, e144989.
61. Hamid, R., Khan, M. A., Ahmad, M., Ahmad, M. M., Abdin, M. Z., Musarrat, J., & Javed, S. (2013). Chitinases: an update. *Journal of Pharmacy and Bioallied Sciences*, 5(1), 21-29.

62. Sandhu, J. S., Sidhu, M. K., & Yadav, I. S. (2017). Control of fungal diseases in agricultural crops by chitinase and glucanase transgenes. In *Sustainable agriculture reviews* (pp. 163-212). Cham: Springer International Publishing.
63. Reicosky, D. C., García-Torres, L., Benítez, J., Martínez-Villela, A., & Holgado-Cabrera, A. (2003). Conservation agriculture. *Climate Change and Terrestrial Carbon Sequestration in Central Asia [Internet]*. Taylor & amp, 199-209.
64. Shokri, N., Robinson, D. A., Afshar, M., Alewell, C., Aminzadeh, M., Arthur, E., ... & Or, D. (2025). Rethinking global soil degradation: Drivers, impacts, and solutions. *Reviews of geophysics*, 63(4), e2025RG000883.
65. Odoh, C. K., Eze, C. N., Obi, C. J., Anyah, F., Egbe, K., Unah, U. V., ... & Adobu, U. S. (2020). Fungal biofertilizers for sustainable agricultural productivity. In *Agriculturally Important Fungi for Sustainable Agriculture: Volume 1: Perspective for Diversity and Crop Productivity* (pp. 199-225). Cham: Springer International Publishing
66. Jie Yin, Jiani Wang, Lu Zhao, Zhongliang Cui, Sheng Yao, Guoxue Li, & Jing Yuan (2026). Compost tea: Preparation, utilization mechanisms, and agricultural applications potential – A comprehensive review, *Environmental Technology & Innovation*, Volume 38.
67. G.Corrado., A.M., M.Z., L.S., & B. Basile (2026), The role and efficacy of compost tea in sustainable fruit tree production: Current scientific knowledge and future directions, *Environmental Technology & Innovation*, Volume 41.
68. Xing, Y., Xie, Y., & Wang, X. (2025). Enhancing soil health through balanced fertilization: a pathway to sustainable agriculture and food security. *Frontiers in Microbiology*, 16, 1536524.
69. N. Jennifer, A. Joseph, (2021) Chapter 18 - Fungi, fungal enzymes and their potential application as biostimulants, *Microbiome Stimulants for Crops*, Woodhead Publishing, Pages 305-314.
70. Garg, S., Kim, M., & Romero-Suarez, D. (2025). Garg, S., Kim, M., & Romero-Suarez, D. (2025). Current advancements in fungal engineering technologies for Sustainable Development Goals. *Trends in Microbiology*, 33(3), 285-301.
71. K.M.H. Nevalainen, (2001) Strain improvement in filamentous fungi-an overview, *Applied Mycology and Biotechnology*, Elsevier, Volume 1, Pages 289-304.
72. Salazar-Cerezo, S., de Vries, R. P., & Garrigues, S. (2023). Strategies for the Development of Industrial Fungal Producing Strains. *Journal of fungi (Basel, Switzerland)*, 9(8), 834.
73. Brahmachari, G., Demain, A. L., & Adrio, J. L. (Eds.). (2016). *Biotechnology of microbial enzymes: production, biocatalysis and Industrial applications*. Academic Press.
74. Garg, S. (2025). The importance of fungal biotechnology for sustainable applications. *Trends in Biotechnology*.
75. Sharma, A., Gupta, G., Ahmad, T., Mansoor, S., & Kaur, B. (2021). Enzyme engineering: current trends and future perspectives. *Food Reviews International*, 37(2), 121-154.
76. Dahiya, P., Panwar, J. S., & Kumar, A. (Eds.). (2024). *Enzyme Biotechnology for Environmental Sustainability*. Elsevier.
77. Banner, A., Toogood, H. S., & Scrutton, N. S. (2021). Consolidated bioprocessing: synthetic biology routes to fuels and fine chemicals. *Microorganisms*, 9(5), 1079.
78. Sharma, V., Nargotra, P., Patel, A. K., & Singhania, R. R. Bioprocessing Approach. *Microbial Fermentation: Innovations, Challenges, and Future Directions*, 139.
79. Dey, P., Rangarajan, V., Singh, J., Nayak, J., & Dilip, K. J. (2022). Current perspective on improved fermentative production and purification of fungal cellulases for successful biorefinery applications: a brief review. *Biomass Conversion and Biorefinery*, 12(3), 967-995.

80. Patel, A. K., Dong, C. D., Chen, C. W., Pandey, A., & Singhania, R. R. (2023). Production, purification, and application of microbial enzymes. In *Biotechnology of microbial enzymes* (pp. 25-57).
81. Ayala, M., Pickard, M. A., & Vazquez-Duhalt, R. (2008). Fungal enzymes for environmental purposes, a molecular biology challenge. *Journal of Molecular Microbiology and Biotechnology*, 15(2-3), 172-180.
82. Kües, U. (2015). Fungal enzymes for environmental management. *Current opinion in biotechnology*, 33, 268-278.
83. Dix, N. J. (Ed.). (2012). *Fungal ecology*. Springer Science & Business Media.
84. Sui, Y., Wisniewski, M., Droby, S., & Liu, J. (2015). Responses of yeast biocontrol agents to environmental stress. *Applied and Environmental Microbiology*, 81(9), 2968-2975.
85. Silva, C., Martins, M., Jing, S., Fu, J., & Cavaco-Paulo, A. (2018). Practical insights on enzyme stabilization. *Critical reviews in biotechnology*, 38(3), 335-350.
86. Khan, M. F. (2025). Fungi for sustainable pharmaceutical remediation: Enzymatic innovations, challenges, and applications—A review. *Processes*, 13(4), 1034.
87. Srivastava, N., Srivastava, M., Mishra, P. K., Gupta, V. K., Molina, G., Rodriguez-Couto, S., ... & Ramteke, P. W. (2018). Applications of fungal cellulases in biofuel production: advances and limitations. *Renewable and Sustainable Energy Reviews*, 82, 2379-2386.
88. Simran, Amin, G., & Kabir, M. G. (2026). Microbial engineering for sustainable microplastic biodegradation: from enzyme redesign to synthetic consortia. *International Microbiology*, 29(1), 23-38.
89. Mehra, M. K., & Avanthi, A. (2024). Challenges and safety regulations for employing fungi in environmental applications. In *Bioprospecting of Multi-tasking Fungi for a Sustainable Environment: Volume I* (pp. 315-341).
90. Magan, N., Fragoeiro, S., & Bastos, C. (2010). Environmental factors and bioremediation of xenobiotics using white rot fungi. *Mycobiology*, 38(4), 238-248.
91. Singh, R. K., Tiwari, M. K., Singh, R., & Lee, J. K. (2013). From protein engineering to immobilization: promising strategies for the upgrade of industrial enzymes. *International journal of molecular sciences*, 14(1), 1232-1277.
92. Maghraby, Y. R., El-Shabasy, R. M., Ibrahim, A. H., & Azzazy, H. M. E. S. (2023). Enzyme immobilization technologies and industrial applications. *ACS omega*, 8(6), 5184-5196.
93. Lynd, L. R., Van Zyl, W. H., McBride, J. E., & Laser, M. (2005). Consolidated bioprocessing of cellulosic biomass: an update. *Current opinion in biotechnology*, 16(5), 577-583.
94. Sharma, B., & Shukla, P. (2020). Designing synthetic microbial communities for effectual bioremediation: A review. *Biocatalysis and Biotransformation*, 38(6), 405-414.
95. Hussain, A., Siddiqu, M., Rehman, U., Ali, H. N., Khan, N. A., Soomru, H. S. P., & Abbas, M. (2025). Applications of Fungal Laccases in the Bioremediation of Industrial Pollutants. *African Journal of Biological Sciences*, 7(197), 2025-196.
96. Sharrock, A., Williams, E., Arcus, V., Mumm, J., & Ackerley, D. (2017). Engineering bacterial nitroreductases for anticancer gene therapy and targeted cell ablation.
97. Gigli, V., Piccinino, D., Avitabile, D., Antiochia, R., Capecchi, E., & Saladino, R. (2022). Laccase mediator cocktail system as a sustainable skin whitening agent for deep eumelanin decolorization. *International journal of molecular sciences*, 23(11), 6238.
98. Andreaus, J., Ximenes Ferreira Filho, E., & Pinto da Silva Bon, E. (2008). Biotechnology of Holocellulose-Degrading Enzymes. *Biocatalysis and bioenergy*, 195-229.

99. Dean, S. N., Turner, K. B., Medintz, I. L., & Walper, S. A. (2017). Targeting and delivery of therapeutic enzymes. *Therapeutic delivery*, 8(7), 577-595.
100. Rana, A., Adhikary, M., Singh, P. K., Das, B. C., & Bhatnagar, S. (2023). "Smart" drug delivery: A window to future of translational medicine. *Frontiers in Chemistry*, 10, 1095598.
101. Naghdi, M., et al. (2017). Bioremediation of petroleum hydrocarbons in soil using fungi: a review. *Environmental Technology Reviews*, 6(3), 139-158. DOI: 10.1080/21622515.2017.1366481
102. Rai, M., et al. (2011). Myconanotechnology: Fungi as an efficient and eco-friendly tool for the synthesis of nanoparticles. *Critical Reviews in Microbiology*, 37(4), 277-286. DOI: 10.3109/1040841X.2011.563342 (Indirect: Fungi use enzymes in nanoparticle synthesis).
103. Santhi, T., et al. (2012). Bioremediation of textile dyes by fungi: a review. *Journal of Microbiology and Biotechnology Research*, 2(4), 613-622.
104. Verma, A., et al. (2020). Fungal enzymes in bioremediation of textile dyes: A review. *Environmental Chemistry Letters*, 18(5), 1481-1494. DOI: 10.1007/s10311-020-01048-0
105. Yadav, A.N., et al. (2023). Fungal diversity in extreme environments: Biotechnological potential and bioremediation strategies. *Extremophiles*, 27(5), 247-269. DOI: 10.1007/s00792-023-01303-x (Enzymes often crucial for survival and function)
106. Zafar, S., & Ali, A. (2019). Fungal enzymes for bioremediation: applications and future perspectives. *Environmental Science and Pollution Research*, 26(31), 31823-31840. DOI: 10.1007/s11356-019-06510-w
107. Singh, H. B., & Vaishnav, A. (Eds.). (2022). *New and future developments in microbial biotechnology and bioengineering: sustainable agriculture: advances in microbe-based biostimulants*. Elsevier.
108. Raghavendra, H., Gulappagouda, R., Basavaraj, D. K., Nayak, G., Singh, P. S., & Sadashiv, S. O. (2025). Emerging Pollutants, Monitoring, and Remediation Techniques: A Sustainable Environmental Applications. In *Biotechnology for Environmental Sustainability* (pp. 187-208). Singapore: Springer Nature Singapore.
109. Jha, H., Mandal, V., Sen, K. K., Tandey, R., Chouhan, K. B. S., & Mehta, R. (2022). Environmental biotechnology: Public perception. In *Emerging Trends in Environmental Biotechnology* (pp. 265-273). CRC Press.
110. Khan, F. (2026). *Sustainable Environmental Biotechnology*. Educohack Press.