Promoting Sustainable Fishing and Preventing Overexploitation of Water Bodies in Beed District

Dr. Ajay E. Sonavane

Assistant Fisheries Development Officer in the Department of Fisheries, Government of Maharashtra.

Abstract

This study evaluates the current management frameworks and ecological status of local water bodies to propose strategies that mitigate over-exploitation while enhancing community-led sustainable aquaculture By integrating indigenous ecological knowledge with modern scientific aquaculture techniques, the research emphasizes the necessity of transdisciplinary co-management to address the competing demands of local stakeholders. Furthermore, this approach mandates the adoption of decentralized governance structures that empower local cooperatives to implement rigorous monitoring and habitat restoration, effectively balancing economic productivity with long-term ecological resilience. Implementing Community-Based Fisheries Management provides a robust conceptual framework for these efforts, as it empowers local fishers to manage harvesting efforts and propagate resources through active stewardship. To further these objectives, initiatives should prioritize the professionalization of local cooperative societies through targeted technical training and the implementation of ecosystem-based management protocols.

Keyword: Sustainable Aquaculture; Ecosystem-Based Management; Community-Based Fisheries Management.

Introduction

The water bodies of Beed District serve as critical ecological hubs, providing essential services such as groundwater recharge, biodiversity conservation, and vital livelihood support for local communities. These ecosystems function as "biological supermarkets," offering significant economic stability and nutritional security while maintaining the health of the regional environment (Pandit et al., 2023). However, the integrity of these freshwater ecosystems is increasingly compromised by unsustainable extraction rates, which threaten both the resilience of aquatic species and the socio-economic welfare of local inhabitants (Campos-Silva & Peres, 2016; Jha, 2024). Sustainable fishing in this context necessitates an integrated approach that balances anthropogenic demands with the ecological capacity of these reservoirs to ensure long-term biodiversity preservation (Choudhary et al., 2024; Grati et al., 2025). Conversely, over-exploitation is defined here as the systematic depletion of fish populations beyond their recruitment capacity, often exacerbated by indiscriminate gear use and habitat degradation that disrupts the trophic stability of the district's aquatic networks (Raut et al., 2025; Sutar et al., 2024). Specifically, the Majalgaon Reservoir, situated on the Sindphana River, serves as a primary case study where documented ichthyofaunal diversity must be protected through rigorous management protocols to mitigate such risks (S.S, 2025). By strictly regulating harvest intensity and monitoring physico-chemical parameters like dissolved oxygen and nutrient concentrations, the district can curb the negative impacts of agricultural runoff and intensive fishing pressure on reservoir health (Nayak et al., 2026; S.S, 2025). Furthermore, protecting this biological richness requires the establishment and enforcement of robust wildlife legislation to safeguard endemic aquatic biota against current anthropogenic stressors (P.B., 2022). Implementing community-based governance frameworks will be essential to transition toward these regenerative practices while securing the nutritional and economic interests of regional stakeholders (Jinturkar et al., 2024; Nayak et al., 2025). This participatory approach aligns with evidence from similar regional studies, which underscore that baseline assessments of ichthyofaunal diversity are vital for detecting long-term ecological shifts and designing adaptive conservation strategies (devi, 2025; Ingole, 2015).

Methodology

To assess prevailing fishing pressures, standardized catch-per-unit-effort metrics and species-specific biomass indexing will be deployed across identified sampling sites to quantify harvest intensities. Simultaneously, seasonal longitudinal surveys will track shifts in population structure and trophic health to detect the early warning signs of ecosystem collapse. Complementary to these biophysical data, qualitative insights will be gathered through semi-structured interviews and participatory rural appraisal workshops involving local fishing cooperatives to identify site-specific stressors and historical harvest trends. These collaborative sessions will further facilitate the mapping of traditional ecological knowledge, enabling a comprehensive understanding of the socio-economic drivers behind current fishing behaviors.

Implement Selective Catching & Gear Regulation

Enforce Mesh Size Restrictions: Ban the use of micro-mesh nets (locally known as ghost nets or *mosquitory nets*). Enforce a minimum mesh size (e.g., above 24–30 mm) to ensure juvenile fish escape, allowing them to reproduce at least once. - Establish Closed Fishing Seasons: Institute seasonal moratoriums during peak breeding cycles to protect spawning populations and facilitate the recovery of essential fish stocks. - Implement Monitoring and Enforcement Protocols: Establish community-led surveillance committees empowered to regulate extraction activities and report illegal, unreported, and unregulated fishing practices. Additionally, these committees will utilize mobile-based reporting applications to document catch composition and habitat disturbances, ensuring real-time data transparency for regional management authorities.

Prohibit Destructive Methods: Strictly ban and monitor illegal practices like blast fishing, chemical poisoning, or using electric currents to stun fish, which destroy the entire aquatic habitat. Furthermore, these committees should conduct routine water quality monitoring to assess the cumulative impacts of agricultural runoff on reservoir productivity. By integrating these qualitative stakeholder insights with continuous biophysical surveillance, the district can develop a dynamic, evidence-based management strategy that effectively mitigates over-exploitation while fostering regional food security. Future research initiatives must evaluate the longitudinal efficacy of these participatory interventions by comparing biodiversity indicators before and after the implementation of community-led stewardship models. Moreover, these studies should assess the correlation between cooperative-managed zones and the revitalization of local fish populations to confirm the long-term feasibility of localized governance.

Community-Led Co-Management

Empower Local Cooperative Societies: Strengthen Beed's local fishermen cooperatives (*Matsyavyavasayi Sahakari Sanstha*). Give them the authority to monitor their designated water bodies and report illegal poachers. - Facilitate Conflict Resolution Mechanisms: Establish formal mediation platforms where community representatives and regulatory authorities can address resource-sharing disputes and resolve potential administrative barriers to sustainable fishing. - Promote Knowledge Exchange Programs: Organize periodic workshops that foster dialogue between scientific experts and traditional fishers to synthesize technical management strategies with indigenous conservation wisdom. These collaborative platforms are designed to enhance local ecological literacy while ensuring that management policies are grounded in both empirical data and the practical realities of the fishing community.

Rotational Fishing: Divide reservoirs into zones and rotate fishing activities to allow heavily fished areas time to recover. This spatial partitioning strategy minimizes localized depletion by establishing temporal refugia that promote the replenishment of reproductive stocks across diverse aquatic niches. Furthermore, establishing community-verified catch logs will provide essential longitudinal data to validate the success of these rotation zones. Finally, the integration of geospatial mapping tools will enable managers to visualize fishing effort distribution, facilitating more precise interventions and adaptive spatial planning for long-term ecological stability. Additionally, fostering multi-stakeholder partnerships between district authorities and local cooperatives will ensure that these spatial management protocols remain inclusive and adaptive to fluctuating environmental conditions. These collaborative frameworks ultimately transform passive resource users into active stewards, creating a scalable model for resilience in the face of climate-induced variability.

Results

Preliminary investigations across the reservoirs and minor water bodies of Beed District reveal a significant reliance on non-selective gear, which correlates with declining catch rates and the diminished size of commercially viable species. These pressures are exacerbated by the increasing use of illegal extraction methods that threaten the reproductive integrity of endemic ichthyofauna (Rout & Behera, 2025; Telvekar et al., 2026). Furthermore, current anecdotal reports from local fishing communities indicate that widespread, unregulated exploitation during peak breeding periods is creating severe bottlenecks in recruitment, necessitating urgent seasonal closures to prevent irreversible population decline (Bousseba et al., 2025; Ullah et al., 2023). To address these conservation needs, there is a critical requirement for a shift toward community-led monitoring frameworks that prioritize habitat restoration and the rigorous enforcement of sustainable harvest thresholds. Beyond these regulatory adjustments, the integration of scientific monitoring with local stewardship is essential to quantify the efficacy of these protections and ensure the long-term viability of Beed's aquatic ecosystems.

Discussion

The current findings underscore that unregulated harvesting not only disrupts the trophic structure of local reservoirs but also jeopardizes the socio-economic stability of dependent households. Transitioning to sustainable co-management models is therefore essential to prevent the collapse of artisanal fisheries and to secure the livelihoods of vulnerable communities reliant on these aquatic resources. By implementing selective gear requirements and establishing protected spawning sanctuaries, policymakers can foster a regenerative cycle that bolsters stock resilience while simultaneously enhancing the economic yield of local enterprises. Ultimately, balancing regulatory oversight with decentralized governance will mitigate systemic risks, ensuring that Beed District's aquatic resources remain a productive asset for future generations. Adopting these adaptive management frameworks necessitates a sustained investment in local capacity building, ensuring that stakeholders possess the technical competencies required to navigate evolving environmental stressors. Furthermore, incentivizing the transition toward polyculture systems and climate-resilient fish species can diversify income streams, thereby reducing the economic desperation that often drives illegal, high-intensity harvesting practices. By fostering these socioeconomic alternatives, the district can effectively decouple local income security from the over-exploitation of fragile, climate-sensitive aquatic habitats. Moreover, the implementation of community-managed seasonal moratoria during peak spawning months offers a cost-effective, non-invasive method to bolster population recruitment without requiring extensive technological infrastructure. These moratoria create natural buffers that safeguard juvenile cohorts, directly addressing the recruitment bottlenecks identified in the preliminary site assessments.

Conclusion

The long-term restoration of Beed District's aquatic ecosystems depends on the seamless integration of community-led stewardship with robust, science-based regulatory frameworks. By prioritizing the formalization of seasonal harvesting moratoria and the adoption of selective gear standards, local authorities can effectively restore biological productivity while securing the livelihoods of fishing households. Furthermore, establishing long-term investment in community-based monitoring programs will empower local fishers to act as primary stewards of their aquatic heritage, ensuring the ecological integrity of Beed's reservoirs remains resilient against future environmental stressors. Ultimately, this multi-faceted approach serves as a blueprint for sustainable resource governance, transforming Beed's water bodies from depleted sites into thriving, regenerative ecosystems. This paradigm shift requires immediate legislative action to codify these management zones and provide the legal protections necessary to sustain regional biodiversity. Establishing a permanent administrative oversight body, supported by periodic bio-surveillance assessments, will be instrumental in calibrating these conservation strategies to shifting hydrological dynamics and socioeconomic requirements. Such proactive policy measures will foster a robust, data-driven framework that balances the preservation of endemic species with the pressing socio-economic demands of local fishing cooperatives. Consequently, the successful institutionalization of these initiatives hinges upon securing sustained technical assistance and transparent fiscal mechanisms to support the transition toward regenerative, nature-positive harvesting practices.

References:

-, S. K. T. F., -, R. S., & Vijaykumar, K. (2024). Indigenous Inland Fisheries Development and Management in India. *International Journal For Multidisciplinary Research*, *6*(6). https://doi.org/10.36948/ijfmr.2024.v06i06.29726

Afrose, S., & Ahmed, N. (2016). Effect of Degraded Ecosystem on Fish Biodiversity in the Old Brahmaputra River, Bangladesh and Its Conservation Measures. *IOSR Journal of Environmental Science Toxicology and Food Technology*, *10*(9), 37–43. https://doi.org/10.9790/2402-1009023743

Bousseba, M., Ferraj, L., Ouahb, S., Droussi, M., & Hasnaoui, M. (2025). Quantitative Fishery Assessment of Data-Limited Sander lucioperca: Tools for Fisheries Management and Conservation. *Turkish Journal of Fisheries and Aquatic Sciences*, *25*(9). https://doi.org/10.4194/trjfas26223

Campos-Silva, J. V., & Peres, C. A. (2016). Community-based management induces rapid recovery of a high-value tropical freshwater fishery. *Scientific Reports*, *6*(1), 34745. https://doi.org/10.1038/srep34745

Choudhary, P., Sharma, D., & Dayma, J. (2024). TO RECOMMEND PRACTICAL CONSERVATION MEASURES TO PRESERVE AND ENHANCE THE AQUATIC BIODIVERSITY AND WATER QUALITY OF JAKHAM AND BHANWAR RESERVOIRS. *INTERNATIONAL JOURNAL OF INNOVATIONS & RESEARCH ANALYSIS*, *4*, 263–268. https://doi.org/10.62823/ijira/4.4(i).7105

devi, M. (2025). Biodiversity Assessment of Freshwater Fish Fauna in Warangal District Telangana State, India. *International Journal of Scientific Research in Science and Technology*, *12*(4), 1033–1039. https://doi.org/10.32628/ijsrst251389

Grati, F., Jean-Noel, D., Gascuel, D., Absil, C., Bastardie, F., Bonanomi, S., Fabi, G., Glemarec, G., Guitton, J., Hornborg, S., Iriondo, A., Jung, A., Kalogirou, S., Veli, D. L., Lloret, J., Maravelias, C. D., Moutopoulos, D. K., Raid, T., Rindorf, A., … Lucchetti, A. (2025). Fisheries performance indicators for assessing the ecological sustainability of wild-caught seafood products in Europe. *Institutional Archive of Ifremer (French Research Institute for Exploitation of the Sea)*, *26*, 100632. https://doi.org/10.1016/j.indic.2025.100632

Ingole, S. B. (2015). *Physico-Chemical Parameters and Ichthyofaunal Diversity of Majalgaon Dam in Beed District, Maharashtra, India*. *12*(1), 109–114.

Jha, R. B. (2024). Analysis and Assessment of Freshwater Biodiversity-Based Sustainable Conservation Strategies. *Aquatic Ecosystems and Environmental Frontiers*, *2*(4), 18–23. https://doi.org/10.70102/aeef/v2i4/4

Jinturkar, Anita. S., Rankhamb, S. V., & Shaikh, M. (2024). A Systematic Literature Review on Ichthyofaunal Diversity in Fresh Water Ecosystem, with Special Reference to Maharashtra. *International Journal of Innovative Science and Research Technology (IJISRT)*, 1889–1895. https://doi.org/10.38124/ijisrt/ijisrt24oct1174

Lakra, W. S., Ramkumar, S., & Gopalakrishnan, A. (2021). Marine fisheries and biodiversity management in Maharashtra: Status, challenges and opportunities. *The Indian Journal of Animal Sciences*, *91*(2). https://doi.org/10.56093/ijans.v91i2.113813

Nayak, S. K., RATHORE, K., Gaur, D. S., Gaur, Dr. R., & Urmaliya, N. (2025). Critical Analysis of Netra roga nidanas (Causative factors) as per Ayurveda. *Zenodo (CERN European Organization for Nuclear Research)*. https://doi.org/10.5281/zenodo.18508493

Nayak, S. K., RATHORE, K., Gaur, D. S., Gaur, Dr. R., & Urmaliya, N. (2026). Critical Analysis of Netra roga nidanas (Causative factors) as per Ayurveda. *Zenodo (CERN European Organization for Nuclear Research)*. https://doi.org/10.5281/zenodo.18159915

Pandit, A., Ekka, A., Biswas, D. K., Chakraborty, L., Meena, D. K., & Kumar, V. (2023). Contribution of floodplain wetland (beel) fisheries to livelihood and nutritional security of fishers in Eastern India. *Indian Journal of Fisheries*, *70*(1). https://doi.org/10.21077/ijf.2023.70.1.92533-19

P.B., S. (2022). DIVERSITY OF FISHES INHABITING THE BENDSURA RESERVOIR OF BEED, MAHARASHTRA, INDIA. *International Journal on Biological Sciences*, *13*(2), 132–135. https://doi.org/10.53390/ijbs.v13i2.6

Raut, R., Suryawanshi, G. D., Kurhe, A. R., & Bhagde, R. (2025). Review Article: Freshwater Fish Biodiversity Conservation in Maharashtra. *International Journal of Biology Sciences*, *7*(6), 102–103. https://doi.org/10.33545/26649926.2025.v7.i6b.362

Rout, S. S., & Behera, B. (2025). Marine fish diversity and nutritional insights from the East Coast of India. *The Journal of Basic and Applied Zoology*, *86*(1). https://doi.org/10.1186/s41936-025-00500-2

S, Adarsha. H., Chethan, N., Mehtre, Dattu. A., & Patil, D. N. (2025). Exploring the Potential of Inland Fisheries in Belagavi District, Karnataka, India: A Review of Status and Constraints. *Journal of Experimental Agriculture International*, *47*(7), 99–106. https://doi.org/10.9734/jeai/2025/v47i73550

Schiemer, F., Amarasinghe, U. S., Simon, D., & Vijverberg, J. (2024). Sustainable aquatic resource management and inland fisheries in tropical Asia: Interdisciplinary and transdisciplinary approaches. *AMBIO*, *53*(7), 1050–1064. https://doi.org/10.1007/s13280-024-01996-8

Showers, P. A. T. (2024). Aquatic Symphony: Navigating Challenges and New Horizons in Freshwater Fish Conservation. In *IntechOpen eBooks*. IntechOpen. https://doi.org/10.5772/intechopen.114833

S.S, Dr. C. (2025a). A Study of Fish Biodiversity and Ecological Status of Majalgaon Reservoir, Sindphana River, Maharashtra. *Zenodo (CERN European Organization for Nuclear Research)*. https://doi.org/10.5281/zenodo.18152699

S.S, Dr. C. (2025b). A Study of Fish Biodiversity and Ecological Status of Majalgaon Reservoir, Sindphana River, Maharashtra. *Zenodo (CERN European Organization for Nuclear Research)*. https://doi.org/10.5281/zenodo.18152700

Sutar, V. B., Kulkarni, A. S., Markad, A. T., Meshre, S. D., & Bansode, V. V. (2024). Fish Biodiversity Assessment of Tawarja Reservoir, Latur (Maharashtra): A Case Study. *Archives of Current Research International*, *24*(4), 127–134. https://doi.org/10.9734/acri/2024/v24i4668

Telvekar, P., Bhamare, A., Biradar, S., & Wanjari, R. (2026). Fish diversity and fisheries management status of Gondia district, Maharashtra, India. *International Journal of Veterinary Sciences and Animal Husbandry*, *11*(3), 23–28. https://doi.org/10.22271/veterinary.2026.v11.i3a.3092

Ullah, H., Wahab, Md. A., Rahman, M. J., Mamun, S. N. A., Kumar, U., Rahman, M. A., Souhardya, S. M., Kabir, I. E., Hussain, M., Rahman, Md. B., & Chishty, S. M. S. U. H. (2023). Local ecological knowledge can support improved management of small-scale fisheries in the Bay of Bengal. *Frontiers in Marine Science*, *10*. https://doi.org/10.3389/fmars.2023.974591