Review - Phytomedicine (Neem)

Pooja Bharti¹, , Pramod Mishra² Sujeet Pratap Singh³, Tarkeshwar Prasad. Shukla⁴

Student, Pharmacy, SCPM College of Pharmacy, Gonda, India ¹
Assistant professor, Pharmacy, SCPM College of Pharmacy, Gonda, India ²
Assistant professor, Pharmacy, SCPM College of Pharmacy, Gonda, India ³
Principal, Pharmacy, SCPM College of Pharmacy, Gonda, India ⁴

Abstract: Global health and medical practice seek to merge alternative medicine with evidence-based medicine for a better understanding of the metabolic process and its effects in the human body. An example is the use of complementary medicine like phytotherapy. Azadirachta indica (Neem), a tree originally from India and Myanmar, called by many "The village pharmacy" or "Divine tree" because of its many health properties. In recent times, Neem-derived extracts have been shown to work from anywhere from insect repellent, to supplements to lower inflammation, diabetic control, and even to combat cancer. Herein, we state the health benefits found in diverse compounds and extracts derived from Neem, highlighting the mechanisms and pathways in which Neem compounds produce their effects, while warning that the improper and unstandardized conditions to produce extracts can lead to health issues, particularly certain compounds might have damaging effects on the liver and kidneys.

Keywords: Phytomedicine, Neem, Mechanism of action of Neem, Application of Neem

INTRODUCTION

Phytomedicine can be defined as the herbal medicine with the rapeutic and healing properties. It came into existence since the advent of human civilization. Sheng Nongs Herbal Book is known as one of the preliminary sources of traditional folk knowledge based on the use of herbs in China and dates back to around 3000 BC. It encompasses the details of almost 365 plants, animals, and minerals that find a place in medication. Our Earth houses approximately 420,000 species of plants; however, there is a lack of appropriate knowledge about them and their varied uses. There are three major areas, namely, food (foodstuffs), medicine (folk and traditional medicines), and research (phytochemical analysis), that predominantly find an immense use of herbal preparations and products and hence can be explored further. Gaining experience from random trials and careful observations from animal studies, people belonging to ancient periods started employing herbs as a therapeutic method against several illnesses. Based on this, the ever so popular Chinese herbal medicine (CHM) as well as Indian herbal medicine, native to and prominently developed in ancient China, Japan, Korea, and India, continue to rule and influence the modern health-care even today. As per the estimate of World Health Organization (WHO), herbal medicines are one of the most sought after primary health-care for around 3.5-4 billion people across the world, and a major portion of traditional medicine involves the plant extract-derived medicines and decoction which may also be termed as the "modern herbal medicine" A herbal medicine or a phytopharmaceutical preparation can be defined as a medicine derived exclusively from a whole plant or parts of plants and manufactured in a crude form or as a purified pharmaceutical formulation. Although with the setting in of the industrial revolution and the advancements in organic chemistry, there was an equivalent increase in the preference for synthetic products as well. However, the WHO emphasizes that between nearly 70% and 95% of the population residing in numerous developing countries still rely more on traditional herbal medicines for their primary medication against diseases (Mohamed et al., 2012). Over the last decade, there has been an enormous rise in the products derived from medicinal plants in terms of interest and use. Our planet is being ruled by plants for over 400 million years now that have successfully survived the test of time even after being challenged time and again by herbivores and microbes. Their defense mechanism is attributed to a range of structurally different secondary metabolites that evolved with time at various stages of development and provide protection against attacks by herbivores, bacteria, fungi, and viruses. Some of these metabolites act as signal compounds that can potentially attract animals that pollinate and disperse the seeds. They additionally act as antioxidants and UV protectants. As far as evolutionary pharmacology is concerned, the secondary metabolites of plant comprise an important collection of bioactive compounds selected and propagated naturally to be used as a remedy against various human infections and health disorders

© 2025, IJSREM | <u>https://ijsrem.com</u> DOI: 10.55041/IJSREM54690 | Page 1



International Journal of Scientific Research in Engineering and Management (IJSREM)

Volume: 09 Issue: 11 | Nov - 2025 SJIF Rating: 8.586 **ISSN: 2582-3930**



Azadirachta indica, commonly known as neem, margosa, nimtree or Indian lilac, is a tree in the mahogany family Meliaceae. It is one of the two species in the genus <u>Azadirachta</u>. It is <u>native</u> to the <u>Indian subcontinent</u> and to parts of <u>Southeast Asia</u>, but is naturalized and grown around the world in <u>tropical</u> and <u>subtropical</u> areas. Its fruits and seeds are the source of <u>neem oil</u>. Nim is a <u>Hindustani</u> noun derived from Sanskrit nimba (निंब).

Kingdom: <u>Plantae</u>

Clade: <u>Tracheophytes</u>

Clade: <u>Angiosperms</u>

Clade: Eudicots

Clade: Rosids

Order: <u>Sapindales</u>

Family: Meliaceae

Genus: Azadirachta

Species: A. indica

Binomial name

Azadirachta indica

The neem tree is a fast-growing <u>tree</u> that can reach a height of 15-20 metres (49–66 ft), and rarely 35-40 m (115-131 ft). It is <u>evergreen</u>, shedding many of its leaves during the dry winter months. The branches are wide and spreading. The fairly dense crown is roundish and may reach a diameter of 20-25 m (66-82 ft). The opposite, <u>pinnate</u> leaves are 20-40 cm (8-16 in) long, with 20 to 30 medium to dark green leaflets about 3-8 cm ($1+\frac{1}{4}-3+\frac{1}{4}$ in) long. The terminal leaflet often is missing. The <u>petioles</u> are short.



The <u>fruit</u> is a smooth (<u>glabrous</u>), olive-like <u>drupe</u> which varies in shape from elongate oval to nearly roundish, and when ripe is 14-28 mm ($\frac{1}{2}-1+\frac{1}{8}$ in) by 10-15 mm ($\frac{3}{8}-\frac{5}{8}$ in). The fruit skin (exocarp) is thin and the bitter-sweet pulp (mesocarp) is

yellowish-white and very fibrous. The mesocarp is 3-5 mm ($\frac{1}{8}-\frac{1}{4}$ in) thick. The white, hard inner shell (endocarp) of the fruit encloses one, rarely two, or three, elongated <u>seeds</u> (kernels) having a brown seed coat.

Phytochemicals

Neem fruit, seeds, leaves, stems, and <u>bark</u> contain diverse <u>phytochemicals</u>, some of which were first discovered in azadirachta seed <u>extracts</u>, such as <u>azadirachtin</u> established in the 1960s as an insect <u>antifeedant</u>, growth disruptor, and insecticide. The yield of azadirachtin from crushing 2 kg of seeds is about 5 g. [19]


© 2025, IJSREM | https://ijsrem.com DOI: 10.55041/IJSREM54690 | Page 2



International Journal of Scientific Research in Engineering and Management (IJSREM)

Volume: 09 Issue: 11 | Nov - 2025 SJIF Rating: 8.586 **ISSN: 2582-3930**

In addition to azadirachtin and related <u>limonoids</u>, the seed oil contains <u>glycerides</u>, diverse <u>polyphenols</u>, nimbolide, <u>triterpenes</u>, and <u>beta-sitosterol</u>. The yellow, bitter oil has a <u>garlic</u>-like odor and contains about 2% of limonoid compounds. The leaves contain <u>quercetin</u>, <u>catechins</u>, <u>carotenes</u>, and <u>vitamin C</u>.



Neem is a <u>tree</u> in the <u>mahogany</u> family <u>Meliaceae</u>. It is native to <u>India</u>, <u>Myanmar</u>, <u>Bangladesh</u>, <u>Sri Lanka</u>, <u>Malaysia</u> and <u>Pakistan</u>. It grows in <u>tropical</u> and semi-tropical regions. It was also the state tree of <u>Hyderabad Deccan</u>.

Neem is a fast-growing <u>tree</u> in India that can reach up to 15–20 m (about 50–65 feet) tall, and sometimes even to 35–40 m (115–131 feet). It is <u>evergreen</u>. It is found in some major parts of Tamil nadu like tuticorin, tirunelyeli, etc.

Uses

Products made from neem have been used in <u>India</u> for over two <u>millennia</u> for their medicinal properties. They are said to be antifungal, <u>antidiabetic</u>, <u>antibacterial</u>, <u>antiviral</u>, <u>contraceptive</u> and <u>sedative</u>. Neem products are also used in selectively controlling <u>pests</u> in plants. Neem is considered a part of <u>Ayurvedic</u> medicine. [11][2][3]

Neem is also known as the village pharmacy. All parts of neem are used for preparing many different medicines, especially for skin disease. [4]


- A compound from the Neem tree can be used as a <u>spermicide</u>.
- Neem oil is used for preparing <u>cosmetics</u> (<u>soap</u> and <u>shampoo</u>, ozone as well as <u>lotions</u> and others), and is useful for skin care such as <u>acne</u> treatment. Neem oil has been used effectively as a <u>mosquito repellent</u>.
- Neem is useful for damaging over 500 types of insects, mites, ticks, and nematodes, by changing the way they grow and act. Neem does not normally kill pests right away, rather it slows their growth and drives them away. As neem products are cheap and not poisonous to animals and friendly insects, they are good for pest control
- In the UK, plant protection products that contain azadirachtin, the active ingredient of neem oil, are illegal. [5]

The plant is well recognized for its medicinal as well as health benefits that it provides to mankind. The presence of certain compounds makes this plant a must-add to your daily routine. It is specifically used for treating outer skin infections like

© 2025, IJSREM | <u>https://ijsrem.com</u> DOI: 10.55041/IJSREM54690 | Page 3

fungi and bacteria. The two medicinal compounds that make Neem essential to be used for treating or preventing fungal infection are nimbidol and gedunin. It possesses the capability to fight and inhibit the growth of fungus spores. [6]

ISSN: 2582-3930



Application:

Neem in combination with aloe vera gel aids in effective recovery from fungal infections as the goodness of aloe vera provides short-term relief from the pain and itchiness caused by the infection.

For external use, wash the infected area with warm neem water which helps in treating fungal infection.

Crush dry neem leaves and make powder.

Mix neem powder with sandalwood water and make a thick paste with rose water.

This mixture is to be applied on the affected area which helps prevent fungal infection.

Note:

Do not apply neem extract directly on infected skin as it may cause itchiness.

Conclusion

Neem is an extraordinarily versatile and ecologically beneficial plant, serving as a natural resource with widespread applications in agriculture, medicine, and the environment. Its rich composition of bioactive compounds, such as azadirachtin, provides antimicrobial, anti-inflammatory, anti-diabetic, and anti-carcinogenic properties. This makes neem invaluable as a sustainable and environmentally friendly alternative to synthetic pesticides and medicines. Continued research into neem's complex chemical interactions and mechanisms will likely unlock even more applications and insights for future human welfare and global ecological health.

References

- Barstow, $M_{\cdot \cdot}$ (2018). "Azadirachta indica". IUCN Red List of Threatened *Deepu*, Species, 2018 e.T61793521A61793525. doi:10.2305/IUCN.UK.2018-1.RLTS.T61793521A61793525.en. Retrieved 19 November 2021.
- 2. "Azadirachta indica A.Juss". Plants of the World Online. Board of Trustees of the Royal Botanic Gardens, Kew. 2017. Retrieved 19 November 2020.
- "Azadirachta indica". Germplasm Resources Information Network. Agricultural Research Service, United States <u>Department of Agriculture</u>. Retrieved 9 June 2017.

© 2025, IJSREM https://ijsrem.com DOI: 10.55041/IJSREM54690 Page 4 IJSREM Le Journal

Oxford University Press.

- 4. Compact Oxford English Dictionary (2013), Neem, page 679, Third Edition 2008 reprinted with corrections 2013,
- 5. <u>Henry Yule</u> and <u>A. C. Burnell</u> (1996), <u>Hobson-Jobson</u>, Neem, page 622, The Anglo-Indian Dictionary, Wordsworth Reference. (This work was first published in 1886)
- 6. Encarta World English Dictionary (1999), Neem, page 1210, St. Martin's Press, New York.
- 7. "Neem | Azadirachta indica". Plants For A Future. 8 January 2018. Retrieved 25 October 2023.
- 8. "Factsheet Azadirachta indica (Neem)". keys.lucidcentral.org. Retrieved 7 August 2024.
- 9. saikia C (2 January 2023). <u>"How to Use Neem- Uses and benefits prakasti.com"</u>. Prakasti. Retrieved 11 January 2023.
- 10. The Tree. National Academies Press (US). 1992.
- 11. "Azadirachta indica A.Juss." The International Plant Names Index. Retrieved 26 March 2023.
- 12. Linnaeus C (1753), "Melia", Species Plantarum, vol. 1, Stockholm: Laurentius Salvius, pp. 384–385, retrieved 26 March 2023
- 13. de Jussieu A (1830). <u>"Mémoire sur le groupe des Méliacées"</u>. Mémoires du Muséum d'histoire naturelle. **19**: 153–304. Retrieved 26 March 2023. <u>p. 221</u>
- 14. "azedarach". Merriam-Webster Dictionary. Retrieved 26 March 2023.
- 15. K. K. Singh, ed. (2009). Neem A Treatise. I.K. International Publishing House, India. p. 3. ISBN 9788189866006.
- 16. <u>Plant Risk Assessment, Neem Tree, Azadirachta indica</u> (PDF). Biosecurity Queensland. 2008. Retrieved 24 January 2014.
- 17. <u>Neem has been declared: what you need to know</u> (PDF), Department of Land Resource Management, 2015, archived from the original (PDF) on 24 March 2015, retrieved 17 March 2015
- 18. <u>Neem Azadirachta indica</u> (PDF), Department of Land Resource Management, 2015, archived from <u>the original</u> (PDF) on 24 March 2015, retrieved 17 March 2015
- 19. "Neem". Drugs.com. 13 August 2020. Retrieved 21 September 2020.
- 20. Anna Horsbrugh Porter (17 April 2006). "Neem: India's tree of life". BBC News.
- 21. "Nimbolide". PubChem, US National Library of Medicine. 6 March 2021. Retrieved 10 March 2021.
- 22. <u>"Different Strokes for Different Folks: A History of the Toothbrush Page 4 Virtual Dental Museum"</u>. dentalmuseum.pacific.edu. Retrieved 8 July 2021.
- 23. Kilani-Morakchi S, Morakchi-Goudjil H, Sifi K (20 July 2021). <u>"Azadirachtin-Based Insecticide: Overview, Risk Assessments, and Future Directions"</u>. Frontiers in Agronomy. 3: 676208. <u>doi:10.3389/fagro.2021.676208</u>.
- 24. "Neem Cake Fertilizer, Uses, Application, Benefits | Agri Farming". www.agrifarming.in. 9 August 2020.
- 25. Minwuyelet A, Yewhalaw D, Aschale Y, et al. (9 June 2025). Gusain P (ed.). "A Global Systematic Review on the Potential of Metal-Based Nanoparticles in the Fight Against Mosquito Vectors". Journal of Tropical Medicine (1). doi:10.1155/jotm/2420073. ISSN 1687-9686. PMC 12170094. PMID 40524903.

© 2025, IJSREM | https://ijsrem.com DOI: 10.55041/IJSREM54690 | Page 5

- 26. Schroeder P (1992). "Carbon storage potential of short rotation tropical tree plantations". Forest Ecology and Management. 50 (1–2): 31–41. <u>Bibcode:1992ForEM..50...31S</u>, <u>doi:10.1016/0378-1127(92)90312-W</u>.
- 27. Puhan, Sukumar, et al. "Mahua (Madhuca indica) seed oil: a source of renewable energy in India." (2005).
- 28. Ramappa KB, Jadhav V, Manjunatha AV (31 May 2022). "A benchmark study on economic impact of Neem Coated Urea on Indian agriculture". Scientific Reports. 12 (9082).
- 29. Heuzé V., Tran G., Archimède H., Bastianelli D., Lebas F., 2015. Neem (Azadirachta indica). Feedipedia, a programme by Institut national de la recherche agronomique, <u>CIRAD</u>, AFZ and <u>FAO</u>. Last updated on 2 October 2015
- 30. Krishnan N, Swetansu Pattnaik, S. A. Deepak, et al. (25 December 2011). "De novo sequencing and assembly of Azadirachta indica fruit transcriptome" (PDF). Current Science. 101 (12): 1553–1561.
- 31. Krishnan N, Swetansu Pattnaik, Prachi Jain, et al. (9 September 2012). "A Draft of the Genome and Four Transcriptomes of a Medicinal and Pesticidal Angiosperm Azadirachta indica". BMC Genomics. 13 (464): 464. doi:10.1186/1471-2164-13-464. PMC 3507787. PMID 22958331.
- 32. Krishnan N, Prachi Jain, Prachi Jain, et al. (20 April 2016). "An Improved Genome Assembly of Azadirachta indica A. Juss". G3: Genes, Genomes, Genetics. 6 (7): 1835–1840. doi:10.1534/g3.116.030056. PMC 4938638. PMID 27172223.
- 33. Narnoliya LK, Rajakani R, Sangwan NS, et al. (2014). "Comparative transcripts profiling of fruit mesocarp and endocarp relevant to secondary metabolism by suppression subtractive hybridization in Azadirachta indica (neem)". Molecular Biology Reports. 41 (5): 3147–3162. doi:10.1007/s11033-014-3174-x. PMID 24477588. S2CID 16605633.
- 34. Rajakani R, Narnoliya L, Sangwan NS, et al. (2014). "Subtractive transcriptomes of fruit and leaf reveal differential representation of transcripts in Azadirachta indica". Tree Genetics & Genomes. 10 (5): 1331–1351. doi:10.1007/s11295-014-0764-7. S2CID 11857916.
- 35. "India wins landmark patent battle". BBC News. 9 March 2005. Retrieved 2 October 2009.
- 36. Sidhu OP, Kumar V, Behl HM (15 January 2003). "Variability in Neem (Azadirachta indica) with Respect to Azadirachtin Content". Journal of Agricultural and Food Chemistry. 51 (4): 910–915. <u>Bibcode: 2003JAFC...51..910S.</u> <u>doi:10.1021/jf025994m. PMID 12568548.</u>
- 37. Prakash G, Bhojwani SS, Srivastava AK (1 August 2002). "Production of azadirachtin from plant tissue culture: State of the art and future prospects". Biotechnology and Bioprocess Engineering. 7 (4): 185–193. doi:10.1007/BF02932968. ISSN 1226-8372. S2CID 85845199.
- 38. Srivastava S, Srivastava AK (17 August 2013). "Production of the Biopesticide Azadirachtin by Hairy Root Cultivation of Azadirachta indica in Liquid-Phase Bioreactors". Applied Biochemistry and Biotechnology. 171 (6): 1351–1361. doi:10.1007/s12010-013-0432-7. ISSN 0273-2289. PMID 23955295. S2CID 36781838.
- 39. Prakash G, Srivastava AK (5 April 2008). "Production of Biopesticides in an in Situ Cell Retention Bioreactor". Applied Biochemistry and Biotechnology. 151 (2–3): 307–318. doi:10.1007/s12010-008-8191-6. ISSN 0273-2289. PMID 18392561. S2CID 35506559

© 2025, IJSREM | https://ijsrem.com DOI: 10.55041/IJSREM54690 | Page 6