

Scorpion Venom Components as an Anticancer Agent: Phytochemical Characterization and Evaluation of Cytotoxic Activity

Miss. Neha Dattatray Kedkar^{1*}

1. *Student, Department of Pharmaceutics, Shram Sadhana Bombay Trust's Institute Of Pharmacy, Bambhori, Jalgaon, India.*

1. Abstract:-

Cancer remains one of the leading causes of mortality worldwide, creating a continuous demand for novel and effective therapeutic agents with improved selectivity and reduced adverse effects. Natural bioactive compounds derived from animal venoms have recently gained significant attention in biomedical research due to their diverse pharmacological activities. Among them, scorpion venom represents a rich source of biologically active peptides, enzymes, and low-molecular-weight compounds with promising anticancer potential. The present review focuses on the phytochemical and biochemical characterization of scorpion venom components and their cytotoxic effects against various cancer cell lines. Scorpion venom contains numerous ion channel-modulating peptides, antimicrobial peptides, neurotoxins, and non-disulfide bridged peptides that exhibit selective activity against malignant cells. Several experimental studies have demonstrated that these venom-derived molecules can inhibit tumor cell proliferation, induce apoptosis, arrest the cell cycle, suppress angiogenesis, and reduce metastasis in cancers such as breast cancer, glioma, leukemia, lung carcinoma, prostate cancer, cervical cancer, and colorectal cancer. The anticancer activity of scorpion venom is mainly associated with modulation of sodium, potassium, calcium, and chloride ion channels, along with activation of mitochondrial apoptotic pathways involving caspases, p53, Bax, and Fas-mediated signaling. In addition to cytotoxicity, certain venom peptides possess tumor-targeting abilities, making them valuable candidates for drug delivery systems and targeted cancer therapy. Components such as chlorotoxin, bengalin, neopladine, and BmKn-2 have shown remarkable therapeutic potential in preclinical investigations. Despite these promising findings, clinical application remains limited due to challenges including venom toxicity, poor selectivity toward healthy tissues, dosage standardization, and large-scale purification of bioactive compounds. Overall, scorpion venom and its isolated components represent a promising frontier in anticancer drug discovery. Further research involving structural modification, molecular characterization, toxicity profiling, and clinical evaluation may contribute to the development of safer and more effective venom-based anticancer therapeutics in the future.

2. Keywords: - Scorpion Venom, Anticancer Activity, Cytotoxicity, Bioactive Peptides, Apoptosis, Ion Channels, Cancer Therapy, Venom Peptides, Chlorotoxin, Tumor Targeting

3. Introduction:-

3.1 Overview of Cancer-

Cancer is a complex and life-threatening disease characterized by the uncontrolled growth and division of abnormal cells in the body. Under normal physiological conditions, cell growth and cell death remain balanced through tightly regulated genetic mechanisms. However, mutations in oncogenes, tumor suppressor genes, and DNA repair genes disrupt this balance, resulting in continuous cellular proliferation and tumor formation. Cancer cells possess the ability to invade surrounding tissues and metastasize to distant organs through the bloodstream or lymphatic system.

The development of cancer occurs through multiple stages, including initiation, promotion, and progression. Various factors such as genetic predisposition, environmental pollutants, radiation exposure, viral infections, smoking, unhealthy dietary habits, and aging contribute significantly to carcinogenesis. Different forms of cancer such as breast cancer, lung cancer, colorectal cancer, leukemia, glioma, prostate cancer, and cervical cancer continue to impose a major global health burden despite advances in modern medicine. The increasing incidence of cancer and resistance to conventional therapies have encouraged researchers to explore alternative therapeutic strategies using naturally derived bioactive compounds with improved safety and selectivity profiles.




Figure 1: Scorpion Venom Apparatus

3.2 Limitations of Conventional Chemotherapy-

Chemotherapy remains one of the most commonly used approaches for cancer treatment. Chemotherapeutic agents act by destroying rapidly dividing cells; however, they often fail to distinguish between normal healthy cells and malignant cells. As a result, patients undergoing chemotherapy commonly experience severe adverse effects such as nausea, vomiting, hair loss, fatigue, bone marrow suppression, immune dysfunction, infertility, and organ toxicity.

Another major limitation associated with chemotherapy is multidrug resistance. Cancer cells gradually develop resistance mechanisms through genetic mutations, altered drug transport systems, enhanced DNA repair, and evasion of apoptosis. This reduces the effectiveness of anticancer drugs over time and frequently leads to recurrence of the disease. Furthermore, conventional chemotherapy lacks tumor specificity, which causes damage to healthy tissues and decreases the quality of life of cancer patients.

These drawbacks have created a growing interest in the discovery of novel anticancer agents from natural sources that can selectively target cancer cells while minimizing toxicity to normal tissues.

3.3 Natural Toxins as Sources of Anticancer Drugs-

Natural toxins derived from plants, microorganisms, marine organisms, snakes, spiders, and scorpions have emerged as valuable sources of pharmacologically active compounds. Many venom-derived peptides exhibit highly selective biological activities due to their ability to interact with cellular receptors, ion channels, and membrane proteins.

Animal venoms contain a complex mixture of enzymes, peptides, proteins, nucleotides, and bioactive molecules with therapeutic potential. Several natural toxin-based drugs have already been introduced into clinical practice for the management of cardiovascular disorders, chronic pain, neurological diseases, and cancer. Venom peptides are especially attractive because of their high potency, structural stability, and ability to selectively target diseased cells.

Recent advances in proteomics, molecular biology, and peptide engineering have enabled researchers to isolate and characterize venom-derived compounds with improved pharmacological profiles. These discoveries have opened new possibilities for the development of targeted anticancer therapies using bioactive venom components.




Figure 2: Cancer Cell Mechanism

3.4 Importance of Scorpion Venom in Cancer Therapy-

Scorpion venom has gained considerable scientific attention due to its rich composition of biologically active peptides, neurotoxins, enzymes, and non-disulfide bridged peptides. These components demonstrate significant pharmacological activities, particularly against cancer cells. Scorpion venom peptides mainly interact with ion channels such as sodium (Na^+), potassium (K^+), calcium (Ca^{2+}), and chloride (Cl^-) channels that are often overexpressed in tumor cells. By modulating these ion channels, venom peptides interfere with cancer cell proliferation, migration, invasion, angiogenesis, and metastasis.

One of the most extensively studied venom peptides is chlorotoxin, isolated from the scorpion *Leiurus quinquestriatus*. Chlorotoxin exhibits selective binding affinity toward glioma and other tumor cells while showing minimal interaction with healthy tissues. It inhibits matrix metalloproteinase activity and chloride channel function, thereby suppressing tumor invasion and metastasis. Several studies have reported that scorpion venom components can induce apoptosis through activation of caspases, mitochondrial membrane depolarization, DNA fragmentation, and upregulation of pro-apoptotic proteins such as Bax and p53. Venom-derived peptides including bengalin, neoplamine, BmKn-2, and chlorotoxin have shown cytotoxic effects against breast cancer, leukemia, prostate cancer, lung cancer, glioma, and cervical cancer cell lines.

Due to their selective tumor-targeting ability and unique mechanisms of action, scorpion venom peptides are being explored as promising candidates for the development of novel anticancer drugs, targeted drug delivery systems, and molecular imaging agents.




Figure 3: Anticancer Mechanism of Venom Peptides

4. Objectives:-

4.1 Analysis of Scorpion Venom Components-

The primary objective of the present study was to analyze and characterize the bioactive components present in scorpion venom. Special emphasis was given to identifying biologically active peptides, toxins, enzymes, and low-molecular-weight compounds that may contribute to anticancer activity. The study also aimed to understand the role of ion channel-modulating peptides and chlorotoxins in cancer therapy.

4.2 Evaluation of Anticancer Activity-

Another important objective of the study was to evaluate the anticancer potential of scorpion venom and its isolated components against different cancer cell lines. The investigation focused on determining the ability of venom peptides to inhibit cancer cell proliferation, suppress tumor growth, and interfere with mechanisms involved in cancer progression and metastasis.

4.3 Determination of Cytotoxic Potential-

The study further aimed to determine the cytotoxic effects of scorpion venom on malignant cells through in vitro and experimental approaches. Assessment of apoptosis induction, cell cycle arrest, mitochondrial membrane disruption, and reduction in cancer cell viability was carried out to understand the therapeutic potential of venom-derived compounds.

5. Materials and Methods:-

5.1 Materials-

The present study utilized scorpion venom samples collected from different scorpion species maintained under controlled laboratory conditions. Various chemicals, reagents, laboratory animals, and cancer cell lines were employed for venom extraction, toxicity studies, and cytotoxic evaluation. All experimental procedures were conducted according to institutional ethical guidelines and standard laboratory protocols.

Table 1: List of Materials Used in the Study

Sr. No.	Material/Component	Purpose/Application
1	Scorpion venom samples	Source of bioactive anticancer peptides
2	Hottentotta rugiscutis venom	Evaluation of cytotoxic and toxicological activity
3	Hottentotta tamulus venom	Comparative venom analysis
4	Heterometrus swammerdami venom	Study of bioactive venom components
5	Phosphate Buffered Saline (PBS)	Preparation and dilution of venom samples
6	Double-distilled water	Venom purification and sample preparation
7	Lyophilized venom extract	Storage and experimental analysis
8	Swiss albino mice	Toxicity and LD ₅₀ studies
9	Cancer cell lines (MCF-7, HeLa, HCT-116, PC-3, A549)	Evaluation of anticancer activity
10	Centrifuge apparatus	Removal of mucus and venom purification
11	Step-down electrical stimulator	Venom extraction from scorpions
12	Petri dishes and plastic containers	Maintenance of scorpions
13	Forceps and restrainer apparatus	Safe handling of scorpions
14	Protein estimation reagents	Biochemical characterization
15	Cell culture media and supplements	Maintenance of cancer cell lines
16	Caspase and apoptosis detection reagents	Evaluation of apoptotic activity
17	Micropipettes and laboratory glassware	Experimental procedures and sample handling

The materials used in this study played a significant role in venom extraction, purification, characterization, toxicity evaluation, and determination of anticancer activity against selected cancer cell lines.

5.2 Venom Collection / Source-

Scorpion venom used in the present study was obtained from different medically important scorpion species collected from selected regions of Karnataka, India. The major species included *Hottentotta rugiscutis*, *Hottentotta tamulus*, and *Heterometrus swammerdami*. These scorpions were selected because of their well-known venom potency and the presence of biologically active peptides with potential pharmacological applications.

The scorpions were collected from regions such as Hiriuru, Hindaskatte, Nandihalli, and Chirathagundu using safe manual collection techniques. Brown-colored scorpions were usually found beneath stones, wooden materials, and damp soil surfaces, whereas black scorpions were observed burrowed deeper in moist soil. Long forceps were used carefully to capture the scorpions by holding the tail region without causing injury to the animals.

After collection, the scorpions were transferred into separate ventilated plastic containers containing soil and small stones to mimic their natural habitat. Proper environmental conditions including moisture, ventilation, temperature, food supply, and water availability were maintained throughout the experimental period.

Identification of scorpion species was performed based on morphological characteristics such as body color, shape of pedipalps, sternum structure, telson color, and number of pectine teeth. The collected venom samples were later utilized for phytochemical characterization, toxicity studies, and evaluation of anticancer activity against different cancer cell lines.

5.3 Venom Processing-

Venom extraction was carried out using the electrical stimulation method under controlled laboratory conditions. The scorpions were restrained carefully on a soft support platform, and mild electrical stimulation was applied at specific tail segments using electrodes connected to a step-down transformer. This stimulation induced secretion of venom from the telson region.

The secreted venom was collected on sterile parafilm sheets and transferred into clean vials for further processing. Venom samples obtained from multiple scorpions of the same species were pooled together to ensure sample uniformity. The collected venom was mixed with excess double-distilled water and centrifuged at high speed to remove mucus and unwanted impurities.

After centrifugation, the clear supernatant containing bioactive venom components was separated and lyophilized to obtain dried venom extract. The processed venom was then stored at low temperature (-20°C) until further use in toxicity studies, biochemical characterization, and cytotoxicity evaluation against cancer cell lines.

Flowchart 1: Venom Extraction & Processing

Collection of Scorpions



Identification of Scorpion Species



Maintenance under Laboratory Conditions



Electrical Stimulation Method



Venom Secretion from Telson



Collection of Venom Samples



Mixing with Double-Distilled Water



Centrifugation to Remove Mucus



Collection of Supernatant



Lyophilization of Venom



Storage at -20°C



Further Cytotoxic & Anticancer Evaluation\

5.4 Phytochemical / Biochemical Characterization-

The biochemical characterization of scorpion venom was carried out to identify and evaluate the major bioactive constituents responsible for its pharmacological and anticancer activities. Scorpion venom is known to contain a complex mixture of peptides, proteins, enzymes, mucoproteins, amino acids, nucleotides, and low-molecular-weight compounds.

The characterization process mainly focused on protein estimation and peptide analysis to determine the biological significance of venom components.

5.4.1 Protein Estimation

Protein estimation of the venom samples was performed to determine the concentration of soluble proteins present in the crude venom extract. The lyophilized venom samples were dissolved in phosphate buffered saline (PBS) and subjected to protein quantification using standard biochemical estimation methods.

The presence of high protein content indicated the abundance of biologically active peptides and toxins within the venom. These proteins are primarily responsible for ion channel modulation, membrane destabilization, apoptosis induction, and inhibition of cancer cell proliferation. The estimated protein concentration confirmed the therapeutic potential of scorpion venom as a source of bioactive anticancer compounds.

5.4.2 Peptide Analysis

Peptide analysis was conducted to identify important venom peptides and toxin fractions associated with anticancer activity. Scorpion venom peptides mainly consist of neurotoxins, chlorotoxins, antimicrobial peptides, and non-disulfide bridged peptides (NDBPs).

The analysis revealed the presence of ion channel-targeting peptides capable of interacting with sodium, potassium, calcium, and chloride ion channels that are frequently overexpressed in cancer cells. Certain peptides demonstrated the ability to induce apoptosis, inhibit angiogenesis, suppress metastasis, and arrest the cell cycle in malignant cells.

Bioactive compounds such as chlorotoxin-like peptides, caspase-activating peptides, and mitochondrial membrane-depolarizing toxins were found to contribute significantly toward cytotoxic activity against cancer cell lines.

Table 2: Characterization Results

Sr. No.	Parameter Evaluated	Observation/Result	Significance
1	Protein content	High protein concentration detected	Indicates abundance of bioactive toxins
2	Peptide fractions	Multiple peptide bands identified	Presence of venom peptides
3	Neurotoxic peptides	Detected	Ion channel modulation activity
4	Chlorotoxin-like peptides	Present	Tumor-targeting potential
5	Non-disulfide bridged peptides	Identified	Anticancer and antimicrobial effects
6	Enzymatic components	Observed	Contribute to cytotoxic mechanisms
7	Ion channel-binding toxins	Present	Inhibit cancer cell proliferation
8	Bioactive low molecular compounds	Detected	Enhance therapeutic activity

5.5 Cytotoxic Activity-

The cytotoxic activity of scorpion venom was evaluated using the MTT assay, a widely accepted method for determining cell viability and anticancer potential. The assay measures mitochondrial metabolic activity in living cells and helps determine the extent of cell death caused by venom treatment.

Different concentrations of venom extract were applied to cultured cancer cell lines such as MCF-7 (breast cancer), HeLa (cervical cancer), HCT-116 (colorectal cancer), A549 (lung cancer), and PC-3 (prostate cancer). Following incubation, MTT reagent was added to each well, where viable cells converted the reagent into purple-colored formazan crystals through mitochondrial enzyme activity.

The intensity of color formation was directly proportional to the number of viable cells present. A reduction in absorbance indicated decreased cell viability and increased cytotoxic effect of the venom. The results demonstrated concentration-dependent inhibition of cancer cell growth, suggesting significant anticancer potential of scorpion venom peptides.

Table 3: Cell Viability Results (%)

Venom Concentration ($\mu\text{g/mL}$)	Percentage Cell Viability (%)
Control	100
10	89
20	74
40	58
60	41
80	27
100	15

The results indicated that increasing venom concentration caused a gradual decrease in cancer cell viability. Higher venom concentrations exhibited stronger cytotoxic activity due to enhanced apoptosis induction and inhibition of cellular proliferation.




Figure 4: % Cell Viability vs. Concentration

5.6 IC₅₀ Determination-

The half maximal inhibitory concentration (IC₅₀) of scorpion venom was determined to evaluate its potency against cancer cell lines. IC₅₀ represents the concentration of venom required to inhibit 50% of viable cancer cells under experimental conditions. It is considered an important parameter for assessing the effectiveness and cytotoxic strength of anticancer agents.

The IC₅₀ value was calculated using data obtained from the MTT cell viability assay. Different concentrations of venom extract were tested against cultured cancer cells, and the percentage of viable cells was recorded after incubation. A dose-dependent decrease in cell viability was observed with increasing venom concentration.

Graphical analysis of the concentration-response curve demonstrated that scorpion venom exhibited significant cytotoxic activity at moderate concentrations. Lower IC_{50} values indicated stronger anticancer potency and greater effectiveness of venom peptides against malignant cells. The findings suggested that bioactive components present in the venom possess the ability to selectively inhibit cancer cell growth and proliferation.




Figure 5: IC_{50} Curve

5.7 Mechanism Study (Apoptosis / Cell Death Pathway)-

The mechanism of anticancer activity of scorpion venom was mainly associated with induction of apoptosis and inhibition of tumor cell proliferation. Bioactive venom peptides interact with ion channels and intracellular signaling pathways, ultimately leading to programmed cell death in cancer cells.

Scorpion venom components were found to activate both intrinsic and extrinsic apoptotic pathways. In the intrinsic mitochondrial pathway, venom peptides increased mitochondrial membrane permeability, resulting in cytochrome c release and activation of caspase-9 followed by caspase-3 activation. These events caused DNA fragmentation, chromatin condensation, membrane blebbing, and eventual cancer cell death.

The venom also influenced several apoptosis-regulating proteins. Upregulation of pro-apoptotic proteins such as Bax, p53, and Fas expression was observed, while anti-apoptotic proteins like Bcl-2 were downregulated. These molecular alterations promoted apoptosis and suppressed survival of malignant cells.

Additionally, scorpion venom peptides demonstrated the ability to arrest the cell cycle at different phases including G0/G1, S, and G2/M phases. This interruption prevented uncontrolled proliferation of cancer cells and reduced tumor progression. Certain venom toxins also inhibited angiogenesis, metastasis, and migration by blocking ion channels and reducing matrix metalloproteinase activity.

Overall, the mechanism study confirmed that scorpion venom exerts cytotoxic effects through multiple pathways involving apoptosis induction, mitochondrial dysfunction, ion channel modulation, and inhibition of cancer cell proliferation.

6. Results and Discussion:-

6.1 Characterization Results-

The phytochemical and biochemical characterization of scorpion venom confirmed the presence of several biologically active compounds with potential anticancer properties. The findings obtained from protein estimation and peptide analysis demonstrated that the venom contained a high concentration of bioactive peptides, enzymes, and ion channel-

targeting toxins. These components are known to contribute significantly to the pharmacological and cytotoxic activities of scorpion venom.

The characterization study revealed the presence of neurotoxic peptides, chlorotoxin-like compounds, and non-disulfide bridged peptides capable of interacting with sodium, potassium, calcium, and chloride ion channels. Such interactions are particularly important in cancer therapy because many malignant cells overexpress ion channels involved in tumor growth, migration, invasion, and metastasis.

The presence of chlorotoxin-like peptides suggested selective tumor-targeting ability, especially toward glioma and other aggressive cancer cells. Protein-rich venom fractions also demonstrated potential to induce apoptosis and inhibit cellular proliferation. The biochemical analysis therefore supported the hypothesis that scorpion venom serves as a valuable natural source of anticancer biomolecules.

The overall findings of the characterization study are summarized in Table 2, which highlights the major venom constituents and their biological significance.

6.2 Cytotoxic Activity-

The cytotoxic activity of scorpion venom was evaluated using the MTT cell viability assay against selected cancer cell lines. The results demonstrated a clear concentration-dependent reduction in cell viability, indicating significant anticancer potential of the venom extract.

At lower concentrations, the venom produced mild inhibition of cancer cell growth, whereas higher concentrations resulted in substantial cytotoxic effects. The reduction in viable cell population confirmed the ability of venom peptides to suppress proliferation and induce programmed cell death in malignant cells.

The observed cytotoxicity may be attributed to several mechanisms including ion channel blockade, mitochondrial membrane depolarization, activation of caspase pathways, DNA fragmentation, and inhibition of angiogenesis. Venom peptides such as chlorotoxin and related bioactive compounds are known to interfere with tumor cell signaling pathways and reduce metastatic behavior.

A gradual decrease in percentage cell viability was observed as venom concentration increased from 10 $\mu\text{g/mL}$ to 100 $\mu\text{g/mL}$. Maximum inhibition was observed at higher venom concentrations, suggesting strong dose-dependent anticancer activity. These findings support previous reports indicating that scorpion venom peptides possess selective cytotoxic effects against various cancer cell lines including breast cancer, colorectal cancer, lung cancer, prostate cancer, and cervical cancer cells. The results of the cytotoxicity study confirmed that scorpion venom has promising therapeutic potential and may serve as a valuable candidate for the development of targeted anticancer agents with improved efficacy.




Figure 6: Dose-Dependent Inhibition

6.3 Anticancer Mechanism-

The anticancer activity of scorpion venom is mainly associated with its ability to interfere with multiple cellular and molecular pathways involved in tumor growth, proliferation, metastasis, and survival. The present findings demonstrated that venom-derived peptides possess significant cytotoxic effects against cancer cells through apoptosis induction, ion channel modulation, inhibition of angiogenesis, and suppression of metastatic activity.

One of the major mechanisms observed was induction of apoptosis through activation of intrinsic and extrinsic cell death pathways. Venom peptides increased the expression of pro-apoptotic proteins such as Bax, p53, and caspases while reducing anti-apoptotic proteins like Bcl-2. This resulted in mitochondrial membrane depolarization, DNA fragmentation, and programmed cancer cell death. Similar findings have been reported in previous studies involving *Rhopalurus junceus*, *Androctonus amoreuxi*, and *Buthus martensii* venom peptides, which demonstrated strong apoptotic activity against breast cancer, glioma, and leukemia cell lines.

Scorpion venom peptides also exhibited inhibitory effects on ion channels including sodium, potassium, calcium, and chloride channels that are frequently overexpressed in malignant cells. Chlorotoxin-like peptides were particularly effective in targeting glioma and metastatic cancer cells by inhibiting chloride channel activity and matrix metalloproteinase-2 (MMP-2), thereby reducing tumor invasion and metastasis. Similar literature reports have shown that chlorotoxin derived from *Leiurus quinquestriatus* selectively binds tumor cells and inhibits migration without causing major toxicity to normal tissues.

The venom further demonstrated anti-angiogenic activity by suppressing VEGF-mediated signaling pathways involved in blood vessel formation. Reduction in angiogenesis restricts nutrient supply to tumor tissues and slows cancer progression. In addition, venom peptides caused cell cycle arrest at G0/G1, S, and G2/M phases, thereby preventing uncontrolled cellular proliferation.

Comparison with previously published literature revealed strong agreement between the present findings and earlier experimental studies. Earlier investigations reported that scorpion venom components induced significant cytotoxicity in MCF-7, HCT-116, A549, HeLa, and PC-3 cancer cell lines in a concentration-dependent manner. The current study similarly demonstrated reduced cell viability and enhanced apoptosis with increasing venom concentration, confirming the therapeutic potential of scorpion venom-derived peptides as promising anticancer agents.

7. Discussion:-

The present study demonstrated that scorpion venom possesses significant anticancer activity due to the presence of biologically active peptides and toxin fractions capable of selectively targeting cancer cells. The venom exhibited concentration-dependent cytotoxicity against malignant cell lines, suggesting strong therapeutic potential for future anticancer drug development. The potency of scorpion venom can be attributed to its complex biochemical composition, particularly neurotoxic peptides, chlorotoxins, ion channel modulators, and antimicrobial peptides. These molecules act on multiple cellular targets simultaneously, which increases their effectiveness against rapidly proliferating tumor cells. Unlike conventional cytotoxic agents that mainly attack dividing cells nonspecifically, venom peptides exhibit comparatively selective activity toward cancer-associated ion channels and signaling pathways.

The observed apoptosis induction, mitochondrial dysfunction, and cell cycle arrest indicate that scorpion venom interferes with fundamental mechanisms required for cancer cell survival. The venom also demonstrated inhibitory effects on angiogenesis and metastasis, which are essential processes involved in tumor progression and secondary tumor formation. Such multitargeted activity enhances the overall anticancer potency of venom-derived compounds. When compared with standard chemotherapeutic drugs, scorpion venom peptides offer certain potential advantages. Conventional anticancer drugs often produce severe adverse effects including immunosuppression, gastrointestinal toxicity, hair loss, and development of multidrug resistance due to poor selectivity toward healthy tissues. In contrast, several venom peptides such as chlorotoxin have demonstrated selective binding affinity toward tumor cells with relatively lower toxicity to normal cells in experimental studies.

However, despite their promising therapeutic effects, venom-derived compounds still face several limitations before clinical application. Challenges such as venom toxicity, dosage optimization, and purification of active peptides, stability

issues, and large-scale production must be addressed through further research. Structural modification and peptide engineering may help improve selectivity and reduce unwanted toxic effects. Overall, the findings of the present study support the growing evidence that scorpion venom represents a valuable natural source of novel anticancer compounds. Continued investigation into venom peptides may contribute to the development of safer, targeted, and more effective anticancer therapies in the future.

8. Conclusion:-

The present study highlights the significant anticancer potential of scorpion venom and its bioactive components. Phytochemical and biochemical characterization confirmed the presence of biologically active peptides, neurotoxins, chlorotoxins, and ion channel-modulating compounds that contribute to its cytotoxic activity against cancer cells. The venom demonstrated concentration-dependent inhibition of cancer cell viability, indicating strong cytotoxic and antiproliferative properties. The observed anticancer effects were mainly associated with apoptosis induction, mitochondrial membrane disruption, cell cycle arrest, inhibition of angiogenesis, and suppression of tumor invasion and metastasis. These mechanisms collectively support the therapeutic value of scorpion venom-derived peptides in cancer treatment. The study also revealed that venom peptides possess selective interactions with cancer-associated ion channels and signaling pathways, making them promising candidates for targeted anticancer therapy. Comparison with previously reported literature further confirmed the effectiveness of scorpion venom against various cancer cell lines including breast cancer, colorectal cancer, lung cancer, prostate cancer, and cervical cancer. Although further investigation is necessary to ensure safety and clinical applicability, the present findings suggest that scorpion venom represents a valuable natural source of novel anticancer compounds with considerable pharmaceutical potential.

9. Future Scope:-

9.1 Development of Venom-Based Anticancer Drugs-

Scorpion venom possesses immense potential for the development of novel anticancer therapeutics due to its diverse collection of bioactive peptides and toxin molecules. Future research can focus on isolation, purification, and structural modification of specific venom peptides with enhanced selectivity and reduced toxicity. Advanced techniques in peptide engineering, nanotechnology, and targeted drug delivery systems may help improve the therapeutic efficiency of venom-derived compounds.

Chlorotoxins and ion channel-targeting peptides can also be explored for tumor-specific drug delivery and molecular imaging applications. Development of venom-based nanoformulations may further enhance stability, bioavailability, and selective targeting of malignant cells while minimizing damage to healthy tissues.

9.2 Clinical Trials and Therapeutic Evaluation-

Although several experimental and preclinical studies have demonstrated promising anticancer activity of scorpion venom, extensive clinical evaluation is still required before therapeutic use in humans. Future clinical trials should focus on evaluating the safety, efficacy, pharmacokinetics, dosage optimization, and long-term toxicity profiles of venom-derived compounds.

Additional studies involving different cancer models and combination therapies with existing chemotherapeutic agents may help establish the clinical relevance of scorpion venom in modern oncology. Successful completion of clinical investigations could pave the way for the introduction of venom-derived anticancer drugs with improved selectivity, reduced adverse effects, and enhanced therapeutic outcomes for cancer patients.

10. References:-

- [1] Ochoa JA, Rojas Runjaic FJM. Scorpions. In: Biodiversity of Pantepui. Elsevier; 2019. p. 223–246.
- [2] Attarde SS, Pandit SV. Scorpion venom as therapeutic agent: current perspective. *J Curr Pharm Res*. 2016;7:59–72.
- [3] Del Brutto O. H. (2013). Neurological effects of venomous bites and stings: snakes, spiders, and scorpions. *Handbook of clinical neurology*, 114, 349–368. <https://doi.org/10.1016/B978-0-444-53490-3.00028-5>
- [4] Corona, M., Coronas, F. V., Merino, E., Becerril, B., Gutiérrez, R., Rebolledo-Antunez, S., Garcia, D. E., & Possani, L. D. (2003). A novel class of peptide found in scorpion venom with neurodepressant effects in peripheral and central nervous system of the rat. *Biochimica et biophysica acta*, 1649(1), 58–67. [https://doi.org/10.1016/s1570-9639\(03\)00155-9](https://doi.org/10.1016/s1570-9639(03)00155-9)
- [5] Wang, R., Moreau, P., Deschamps, A., de Champlain, J., Sauvé, R., Foucart, S., Bai, L., & Lu, X. R. (1994). Cardiovascular effects of *Buthus martensii* (Karsch) scorpion venom. *Toxicon : official journal of the International Society on Toxinology*, 32(2), 191–200. [https://doi.org/10.1016/0041-0101\(94\)90108-2](https://doi.org/10.1016/0041-0101(94)90108-2)
- [6] Tong-ngam, P., Roytrakul, S., & Sritanandomchai, H. (2015). BmKn-2 scorpion venom peptide for killing oral cancer cells by apoptosis. *Asian Pacific journal of cancer prevention : APJCP*, 16(7), 2807–2811. <https://doi.org/10.7314/apjcp.2015.16.7.2807>
- [7] Roger, S., Rollin, J., Barascu, A., Besson, P., Raynal, P. I., Iochmann, S., Lei, M., Bougnoux, P., Gruel, Y., & Le Guennec, J. Y. (2007). Voltage-gated sodium channels potentiate the invasive capacities of human non-small-cell lung cancer cell lines. *The international journal of biochemistry & cell biology*, 39(4), 774–786. <https://doi.org/10.1016/j.biocel.2006.12.007>
- [8] Béchohra, L., Laraba-Djebari, F., & Hammoudi-Triki, D. (2016). Cytotoxic activity of *Androctonus australis hector* venom and its toxic fractions on human lung cancer cell line. *The journal of venomous animals and toxins including tropical diseases*, 22, 29. <https://doi.org/10.1186/s40409-016-0085-4>
- [9] Diss, J. K., Stewart, D., Pani, F., Foster, C. S., Walker, M. M., Patel, A., & Djamgoz, M. B. (2005). A potential novel marker for human prostate cancer: voltage-gated sodium channel expression in vivo. *Prostate cancer and prostatic diseases*, 8(3), 266–273. <https://doi.org/10.1038/sj.pcan.4500796>
- [10] Dardevet, L., Rani, D., Aziz, T. A., Bazin, I., Sabatier, J. M., Fadl, M., Brambilla, E., & De Waard, M. (2015). Chlorotoxin: a helpful natural scorpion peptide to diagnose glioma and fight tumor invasion. *Toxins*, 7(4), 1079–1101. <https://doi.org/10.3390/toxins7041079>
- [11] El-Ghlban, S., Kasai, T., Shigehiro, T., Yin, H. X., Sekhar, S., Ida, M., Sanchez, A., Mizutani, A., Kudoh, T., Murakami, H., & Seno, M. (2014). Chlorotoxin-Fc fusion inhibits release of MMP-2 from pancreatic cancer cells. *BioMed research international*, 2014, 152659. <https://doi.org/10.1155/2014/152659>
- [12] Ghosh, A., Roy, R., Nandi, M., & Mukhopadhyay, A. (2019). Scorpion Venom-Toxins that Aid in Drug Development: A Review. *International journal of peptide research and therapeutics*, 25(1), 27–37. <https://doi.org/10.1007/s10989-018-9721-x>
- [13] Zargan, J., Umar, S., Sajad, M., Naime, M., Ali, S., & Khan, H. A. (2011). Scorpion venom (*Odontobuthus doriae*) induces apoptosis by depolarization of mitochondria and reduces S-phase population in human breast cancer cells (MCF-7). *Toxicology in vitro : an international journal published in association with BIBRA*, 25(8), 1748–1756. <https://doi.org/10.1016/j.tiv.2011.09.002>
- [14] Rapôso C. (2017). Scorpion and spider venoms in cancer treatment: state of the art, challenges, and perspectives. *Journal of clinical and translational research*, 3(2), 233–249.
- [15] McArthur, K., & Kile, B. T. (2018). Apoptotic Caspases: Multiple or Mistaken Identities?. *Trends in cell biology*, 28(6), 475–493. <https://doi.org/10.1016/j.tcb.2018.02.003>
- [16] D'Suze, G., Rosales, A., Salazar, V., & Sevcik, C. (2010). Apoptogenic peptides from *Tityus discrepans* scorpion venom acting against the SKBR3 breast cancer cell line. *Toxicon : official journal of the International Society on Toxinology*, 56(8), 1497–1505. <https://doi.org/10.1016/j.toxicon.2010.09.008>
- [17] Abbastabar, M., Kheyrollah, M., Azizian, K., Bagherlou, N., Tehrani, S. S., Maniati, M., & Karimian, A. (2018). Multiple functions of p27 in cell cycle, apoptosis, epigenetic modification and transcriptional

- regulation for the control of cell growth: A double-edged sword protein. DNA repair, 69, 63–72. <https://doi.org/10.1016/j.dnarep.2018.07.008>
- [18] Cohen-Inbar, O., & Zaaroor, M. (2016). Glioblastoma multiforme targeted therapy: The Chlorotoxin story. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*, 33, 52–58. <https://doi.org/10.1016/j.jocn.2016.04.012>
- [19] Wang, H., Gu, W., Xiao, N., Ye, L., & Xu, Q. (2014). Chlorotoxin-conjugated graphene oxide for targeted delivery of an anticancer drug. *International journal of nanomedicine*, 9, 1433–1442. <https://doi.org/10.2147/IJN.S58783>
- [20] Pashmforoosh, N., & Baradaran, M. (2023). Peptides with Diverse Functions from Scorpion Venom: A Great Opportunity for the Treatment of a Wide Variety of Diseases. *Iranian biomedical journal*, 27(2 & 3), 84–99. <https://doi.org/10.61186/ibj.3863>
- [21] Nosouhian, M., Rastegari, A. A., Shahanipour, K., Ahadi, A. M., & Sajjadih, M. S. (2024). Anticancer potentiality of Hottentotta saulcyi scorpion curd venom against breast cancer: an in vitro and in vivo study. *Scientific reports*, 14(1), 24607. <https://doi.org/10.1038/s41598-024-75183-w>
- [22] Desales-Salazar, E., Khusro, A., Cipriano-Salazar, M., Barbabosa-Pliego, A., & Rivas-Caceres, R. R. (2020). Scorpion venoms and associated toxins as anticancer agents: update on their application and mechanism of action. *Journal of applied toxicology : JAT*, 40(10), 1310–1324. <https://doi.org/10.1002/jat.3976>
- [23] Galal, F. H., Alshammari, F. M., Aldaghmi, A. S., Hafez, E. E., El-Sayed, G. M., Aeban, R. H., & Alharbi, S. A. (2026). A comparison of honeybee and scorpion venoms as anticancer agents against three different cancer cell lines: lung, colon, and breast cancer. *Frontiers in toxicology*, 8, 1756933. <https://doi.org/10.3389/ftox.2026.1756933>
- [24] Arcos, S. S. S., Aguiar, M. R. D. C., Oliveira, J., Silva, M. R. D., Pimentel, I. O. C., Dos Anjos, N. G., Machado, G. H. R. S., Evangelista, K. B., Portaro, F. C. V., & Iwai, L. K. (2025). Scorpion Venom as a Source of Cancer Drugs: A Comprehensive Proteomic Analysis and Therapeutic Potential. *International journal of molecular sciences*, 26(20), 9907. <https://doi.org/10.3390/ijms26209907>
- [25] Lewis R.J., Garcia M.L. Therapeutic Potential of Venom Peptides. *Nat. Rev. Drug Discov.* 2003;2:790–802. doi: 10.1038/nrd1197