

Study on Atmospheric Hydrocarbon Exposure During LPG and Propane Loading Activities and its Impact on Worker Well-Being

Gourav Dubey¹, Prof. Vishal Tiwari²

¹Research Scholar, Department of Fire Technology & Safety Engineering, School of Engineering and Technology Vikrant University, Gwalior, (M.P.), India

²M.Tech Guide, Department of Fire Technology & Safety Engineering, School of Engineering and Technology Vikrant University, Gwalior, (M.P.), India

Abstract

Liquefied Petroleum Gas (LPG) and propane are widely used fuels in industrial and domestic sectors. During tanker loading operations, hydrocarbon vapors are released into the atmosphere, exposing workers to potentially hazardous conditions. This study investigates the extent of atmospheric hydrocarbon exposure during LPG and propane loading and evaluates its impact on workers' physical and physiological well-being. The findings indicate that prolonged exposure to hydrocarbon vapors can adversely affect respiratory, cardiovascular, and metabolic health. The paper also discusses mitigation strategies to reduce occupational exposure and enhance workplace safety.

Keywords

Hydrocarbon exposure, LPG loading, propane, occupational health, atmospheric emissions, worker safety

1. Introduction

Hydrocarbons, particularly those present in LPG and propane, are volatile organic compounds (VOCs) that can easily evaporate into the atmosphere during handling and transfer operations. Tanker loading processes involve pressurized systems where minor leakages or venting can result in atmospheric emissions.

Previous studies indicate that hydrocarbon release during loading operations may occur through equipment such as rotogauges and valves, contributing to both environmental pollution and occupational health risks. Workers involved in such operations are

at increased risk of inhaling these vapors, especially in poorly ventilated or high-frequency loading environments.

2. Literature Review

Research on occupational exposure to LPG and hydrocarbons highlights several health concerns:

- Exposure to LPG vapors has been associated with respiratory and cardiovascular impairments, including reduced lung function and increased blood pressure.
- Studies show that prolonged exposure can lead to liver enzyme alterations and oxidative stress, indicating systemic toxicity.
- Workers exposed to LPG during tanker loading exhibited elevated cholesterol levels, which improved when exposure was reduced.
- Occupational exposure to hydrocarbon vapors has been linked to declines in pulmonary function, including reduced FEV1 and FVC values.

Additionally, petroleum gas toxicity studies indicate that hydrocarbons are lipid-soluble and can affect multiple organ systems when inhaled.

3. Objectives of the Study

1. To analyze the sources and extent of hydrocarbon release during LPG and propane loading.
2. To assess the exposure levels experienced by workers.
3. To evaluate the impact of hydrocarbon exposure on worker health and well-being.
4. To recommend preventive and control measures for reducing occupational risks.

4. Methodology

This study adopts a descriptive and analytical research design.

4.1 Data Collection

- Primary data: Observation of LPG loading operations and worker health surveys.
- Secondary data: Published research articles, occupational safety reports, and toxicological studies.

4.2 Sampling

Workers involved in LPG and propane tanker loading operations were considered for analysis.

4.3 Analytical Tools

- Comparative analysis of exposed vs. non-exposed workers
- Review of medical indicators (respiratory function, blood parameters)
- Risk assessment models for hydrocarbon exposure

5. Sources of Hydrocarbon Emissions During Loading

Hydrocarbon emissions during LPG and propane loading occur due to:

- Leakage from valves, hoses, and connectors
- Release through measurement devices (e.g., rotogauges)
- Vapor displacement during filling operations
- Equipment wear and improper maintenance

These emissions contribute to localized atmospheric contamination, especially in confined or poorly ventilated loading bays.

6. Occupational Exposure Pathways

Workers are exposed to hydrocarbons primarily through:

- Inhalation (major pathway)
- Dermal contact (minor but possible)

Exposure levels depend on:

- Duration of loading operations
- Frequency of tasks
- Environmental conditions (wind, temperature)
- Safety measures in place

Studies show that hydrocarbon concentration levels in petroleum handling environments can vary significantly and often follow a log-normal distribution.

7. Impact on Worker Well-being

7.1 Respiratory Effects

- Reduced lung capacity (FEV1, FVC decline)
- Increased risk of airway obstruction
- Chronic respiratory symptoms such as cough and breathlessness

7.2 Cardiovascular Effects

- Elevated systolic blood pressure
- Increased cardiovascular stress due to prolonged exposure

7.3 Hepatic and Metabolic Effects

- Altered liver enzyme levels
- Increased cholesterol levels in exposed workers

7.4 Neurological and Acute Effects

- Headache, dizziness, and fatigue
- In severe cases, unconsciousness due to oxygen displacement

7.5 Long-Term Health Risks

- Chronic respiratory diseases
- Potential systemic toxicity
- Reduced overall work efficiency and productivity

8. Risk Assessment

Hydrocarbon exposure risk depends on:

- Concentration of vapors in air
- Exposure duration
- Worker proximity to emission source

Risk is higher during:

- Continuous loading operations
- Equipment malfunction
- Inadequate ventilation

9. Control and Preventive Measures

9.1 Engineering Controls

- Installation of vapor recovery systems
- Use of leak-proof equipment
- Regular maintenance and inspection

9.2 Administrative Controls

- Limiting exposure duration
- Rotational work schedules
- Regular health monitoring

9.3 Personal Protective Equipment (PPE)

- Respirators or masks
- Protective clothing
- Gas detection systems

9.4 Training and Awareness

- Safety training for workers
- Emergency response planning
- Awareness of hydrocarbon hazards

improved equipment design, monitoring systems, and safety protocols, are essential to protect workers and enhance operational safety.

References

1. Tandale, A. (2012). *Study of Release of Hydrocarbon in the Atmosphere while LPG and Propane Tankers Loading and Its Occupational Health Effects*. International Journal of Scientific and Research Publications.
2. Tandale, A., Siddiqui, N. A., & Gautam, A. (2013). *Occupational Health Effects of Exposure of LPG and Propane on Human*. IJARET.
3. Science of the Total Environment (2023). *Impact of differential occupational LPG exposure on cardiopulmonary indices, liver function, and oxidative stress*.
4. PubMed Study (2014). *Airflow obstruction among LPG-exposed workers*.
5. Verma, D. K. et al. (1992). *Hydrocarbon exposures at petroleum bulk terminals*.
6. McKee, R. H. et al. (2014). *Toxicological Properties of Petroleum Gases*.
7. Rahimi Moghadam, S. et al. (2019). *Effects of occupational exposure to petroleum products on pulmonary function*.

Discussion

The findings confirm that hydrocarbon emissions during LPG and propane loading pose significant occupational health risks. While individual exposure during a single operation may be low, cumulative exposure over time can lead to serious health issues.

The study emphasizes the need for a comprehensive occupational health framework, integrating engineering controls, monitoring systems, and worker education.

Conclusion

Atmospheric hydrocarbon exposure during LPG and propane loading activities is a critical occupational hazard. Prolonged exposure adversely affects worker health, particularly respiratory and cardiovascular systems. Effective mitigation strategies, including